

原著論文（相馬正義）

1. Honda M, Watanabe H, Inoue T, Soma M, Manabe R, Izumi Y: Study on the hypotensive mechanism of angiotensin I converting enzyme inhibitor (SQ 14,225) in the acute phase. The Nihon University Journal of Medicine 24:1982.3
2. Soma M, Honda M, Hatano M: Effects of angiotensin II and angiotensin III on prostacyclin release in man. Japanese Journal of Nephrology 25:1093-1101, 1983.9
3. Honda M, Tsuchiya M, Manabe R, Watanabe Y, Inoue T, Nagashima Y, Izumi Y, Soma M, Watanabe H, Shiratsuchi T, Ueda Y, Fukuda N, Minato M, Watanabe M, Hatano M: Effects of prostacyclin on the responses of blood pressure and several hormones in normotensive subjects. Japanese Journal of Nephrology 26:1235-1241, 1984.9
4. 高山亮子、本多正信、渡辺吉康、福田昇、永嶋康夫、井上毅、湊昌也、渡辺昌司、白土寿、泉洋一、相馬正義、波多野道信：腎機能障害を合併した高血圧患者の腎機能に及ぼす Converting enzyme inhibitor (CEI) 長期投与の影響。新薬と臨床 34:pp.425-431,1985.3
5. Soma M, Manku MS, Jenkins DK and Horrobin DF. Prostaglandins and thromboxane outflow from the perfused mesenteric vascular bed in spontaneously hypertensive rats. Prostaglandins 29:323-333, 1985.7
6. Soma M, Manku MS and Horrobin DF. The effects of hydrogenated coconut oil, safflower oil, and evening primrose oil on development of hypertension and sodium handling in spontaneously hypertensive rats. Canadian Journal of Physiology and Pharmacology 63:325-330, 1985.10
7. Soma M, Manku MS, Jenkins DK, Huang Y-S and Horrobin DF. Effects of dietary saturated, n-6 and n-3 polyunsaturated fats on blood pressure and prostaglandins outflow from perfused mesenteric vascular bed in rats. Prostaglandins 30:969-980, 1985.12
8. Cunnane S.C., Soma M, McAdoo KR and Horrobin DF. Magnesium deficiency in the rat

increased tissue levels of docosahexaenoic acid. *Journal of Nutrition* 115:1498-1503, 1985.12

9. 本多正信、高山亮子、福田昇、湊昌也、渡辺昌司、永嶋康夫、相馬正義、白土寿、泉洋一、波多野道信：Ca拮抗薬によるヒト血中および尿中 Prostaglandins と Thromboxane B2 の動態について。日本腎臓学会雑誌 28:799-806, 1986.6
10. Soma M. Effects of cholesterol feeding on platelet thromboxane synthesis and vascular prostaglandins in rats. *Progress in Lipid Research* 25:205-206, 1986.7
11. Soma M. Fatty acid metabolism and PGI₂ synthesis in mesenteric arterial beds in spontaneously hypertensive rats. *Progress in Lipid Research* 25:323-326, 1986.7
12. Manku M.S, Soma M and Horrobin DF. Effects of feeding gamma-linolenic acid on mesenteric and urinary prostaglandin outflow in guinea-pig. *Progress in Lipid Research* 25:309-310, 1986.7
13. Fujii K, Soma M, Huang YS, Manku MS and Horrobin DF: Increased release of prostaglandins from the mesenteric vascular bed in streptozotocin diabetes. *Progress in Lipid Research* 25:499-501, 1986.7
14. Fujii K, Soma M, Huang YS, Manku MS and Horrobin DF: Effect of glucose and insulin on release of prostaglandins from the mesenteric vascular bed in control and streptozotocin diabetic animal. *Progress in Lipid Research* 25:503-505, 1986
15. 相馬正義、本多正信、湊昌也、波多野道信：Nicardipine Hydrochloride (Perdipine) の肝血流に及ぼす影響について。基礎と臨床 20:pp4907-4909,1986.9
16. Fujii K, Soma M, Huang YS, Manku MS and Horrobin DF: Increased release of prostaglandins from the mesenteric vascular bed of diabetic animals: the effects of glucose and insulin. *Prostaglandins Leukot Med* 24:151-161, 1986.10
17. 小倉敏嗣、佐藤光重、相馬正義、斎藤友昭：心不全症例に対する urinastatin の使用

経験。臨床と研究 64:319-323,1987.9

18. Soma M, SC. Cunnane, DF. Horrobin, MS. Manku, M. Honda, and M. Hatano: Effects of low magnesium diet on the vascular prostaglandin and fatty acid metabolism in rats. Prostaglandins 36:pp431-441, 1988.4
19. 本多正信、湊昌也、渡部昌司、福田昇、相馬正義、永嶋康夫、渡辺吉康、泉洋一、白土寿、波多野道信：高血圧自然発症ラット (SHR) における新しい Ca 拮抗薬 Franidipine (CV-4093 2HCl) の血圧および内分泌系因子に与える効果について。日本脈管学会雑誌 28:1075-1081, 1988.12
20. 相馬正義、本多正信、永嶋康夫、渡部昌司、湊昌也、福田昇、波多野道信：非観血的携帯型自動血圧計による降圧剤の評価-Indapamide について-。基礎と臨床 22:291-294, 1988.12
21. 相馬正義、泉洋一、渡辺吉康、福田昇、波多野道信：血管壁 Eicosanoids 産生に及ぼす Mg 欠乏の影響について。マグネシウム 8:pp.25-28, 1989.1
22. 本多正信、湊昌也、高山亮子、渡部昌司、福田昇、相馬正義、永嶋康夫、泉洋一、渡辺吉康、白土寿、波多野道信、栗本文彦、桜井兵一郎、有賀とし弥：19-hydroxyandrostenedione の昇圧機序についての研究。日本内分泌学会雑誌 65:18-31, 1989
23. 湊昌也、本多正信、相馬正義、波多野道信：C19 ステロイドの昇圧機序について-外因性昇圧物質に対する昇圧反応。日大医学雑誌 48:pp581-585, 1989
24. 渡部昌司、渡部昌司、相馬正義、湊昌也、福田昇、渡辺吉康、永嶋康夫、泉洋一、波多野道信：非観血的携帯型自動血圧計による降圧効果の評価-カプトプリル 25mg 1日2回投与について-。Progress in Medicine 9:pp.1656-1658, 1989.5
25. 相馬正義、泉洋一、渡辺吉康、福田昇、渡辺昌司、波多野道信：非観血的携帯型自動血圧計による降圧剤の評価-持効性 Nifedipine について-。診療と新薬 27:1729-1732, 1990.

26. 渡辺吉康、泉洋一、相馬正義、福田昇、渡辺昌司、波多野道信：非観血的携帯型自動血圧計による降圧薬の評価-塩酸アロチノロールの血圧日内変動に及ぼす効果-。新薬と臨床 40:pp151-154, 1990.1
27. Watanabe M, M. Honda, M. Minato, N. Fukuda, M. Soma, Y. Watanabe, Y. Izumi, M. Hatano: Measurement of intracellular free calcium in human platelets. Nihon University Journal of Medicine 32:123-129, 1990.
28. Fukuda N, Y. Izumi, M. Soma, Y. Watanabe, M. Watanabe, M. Hatano, I. Sakuma, H. Yasuda: L-NG-Monomethyl arginine inhibits the vasodilating effects of low dose of endothelin-3 on rat mesenteric arteries. Biochemical and Biophysical Research Communications 167:739-745, 1990.
29. Fukuda N, M. Honda, M. Minato, M. Soma, Y. Izumi, and M. Hatano: Adenylate cyclase activities of vascular smooth muscle in early and established DOCA/Salt hypertensive rat. Japanese Circulation Journal 54:pp.82-88, 1990.
30. Izumi Y, Honda M, Fukuda N, Watanabe M, Watanabe Y, Soma M, Hatano M: Effect of arterial natriuretic peptide on adrenal renin and aldosterone. Endocrinologica Japonica 37:pp.47-52, 1990.
31. Watanabe M, Honda M, Soma M, Minato M, Fukuda N, Watanabe Y, Izumi Y, Hatano M: Preventive effect of angiotensin I on weight reduction in the adrenal glands of DOCA/Salt hypertensive rats. Endocrinologica Japonica 37:pp.521-528, 1990.
32. Fukuda N, Izumi Y, Soma M, Watanabe Y, Watanabe M, Hatano M, Sakuma I, Yasuda H: Effects of Indomethacin, Endothelium-Denudation, Methylene Blue and L-NG-Monomethyl arginine on the Vasoactive Effects of Endothelin-3. Japanese Journal of Pharmacol 55:375-380, 1991.9
33. Fukuda N, Soma M, Izumi Y, Minato M, Watanabe Y, Watanabe M, Hatano M: Low doses of endothelin-3 elicit endothelium dependent vasodilation which accompanies with

- elevation of cyclic GMP. Japanese Circulation Journal 55:pp.617-622, 1991.6
34. Fukuda N, Izumi Y, Minato M, Soma M, Watanabe Y, Watanabe M, and Hatano M: Cyclic GMP formation of resistance vessels in the development of hypertension in spontaneously hypertensive rats. Japanese Circulation Journal 55:721-726, 1991.7
 35. Watanabe M, Izumi Y, Soma M, Watanabe Y, Fukuda N, Abe Y, Ito M, Kageura H, Nakayama T, Hatano M: Effect of endothelin-3 (ET-3) on renal function in rat perfused kidney. Endocrinologica Japonica 38:435-440, 1991.8
 36. Fukuda N, Izumi Y, Soma M, Watanabe Y, Watanabe M, and Hatano M: Effects of L-NG-Monomethyl arginine on the cyclic GMP formation in rat mesenteric arteries. Japanese Journal of Pharmacology 58:55-60, 1992.
 37. 中山智祥、相馬正義、影浦博信、阿部好伸、伊藤みき、渡辺昌司、渡辺吉康、泉洋一、波多野道信：Sipple 症候群における高血圧型と尿中カテコラミンについて。日本内分泌学会雑誌 68:pp.39-50, 1992.1
 38. 蔵本築、相馬正義、他 28 名：塩酸デラプリルの QOL に対する影響の検討。Geriatric Medicine 31:559-568, 1993.4
 39. Nakayama T, Imai S, Soma M, Izumi Y, Kanmatsuse K: Compensatory adrenal growth and steroidogenesis after unilateral adrenalectomy. Endocrine Journal 40:pp523-527, 1993.10
 40. 天野定雄、黒須康彦、秦怜志、柴田昌彦、三宅洋、滝沢秀博、宗像敬明、相馬正義、上松瀬勝男：のう胞性甲状腺腫に対するエタノール注入療法。日大医学雑誌 53:pp529-532, 1994.6
 41. Ito M, Fukuda N, Izumi Y, Watanabe Y, Watanabe M. Soma M, Kammatsuse K: Enhancement effect of carteorol on the clonidine-induced vasodilation of rat mesenteric arteries. Pharmacological Research 31:319-323, 1995.5
 42. Ito M. Soma M, Kammatsuse K: Effects of long-term treatment with angiotensin converting

- enzyme inhibitor on angiotensin II and prostacyclin release from mesenteric arteries in spontaneously hypertensive rats. *Prostaglandin Leukotrien and EFAs*, 53:359-363, 1995.11
43. Nakayama T, Soma M, Izumi Y, Kanmatsuse K, Esumi M: CA repeat polymorphism of the endothelial nitric oxide synthase gene in Japanese. *Human Heredity* 45:301-302, 1995.8
44. 中山智祥、田沼美昭、相馬正義、上松瀬勝男、大和田操、中林啓記、江角真理子：インスリン非依存性糖尿病と筋型グリコーゲン合成酵素遺伝子マイクロサテライト多型との関連。日本内分泌学会誌、71:597-604, 1995.
45. Fukuda N, Kubo A, Izumi Y, Soma M and Kanmatsuse K: Characteristics and expression of transforming growth factor- β receptor subtypes on vascular smooth muscle cells from spontaneously hypertensive rats. *J Hypertens* 13:831-837, 1995.10
46. Fukuda N, Kubo A, Izumi Y, Soma M, Kanmatsuse K: Distinct expression of transforming growth factor- β receptor subtypes on vascular smooth muscle cells from spontaneously hypertensive rats and Wistar-Kyoto rats. *Clin Experi Pharmacol Physiol. suppl1*, S120-122, 1995.
47. 國本昌子、相馬正義、泉洋一、上松瀬勝男：ラット腸管膜抵抗血管に対する nifedipine の血管拡張作用の検討。 *Therapeutic Research* 16:2689-2696, 1995.8
48. Soma M, Izumi Y, Watanabe Y, Kanmatsuse K: A nitric oxide synthesis inhibitor decreased prostaglandin production in rat mesenteric vasculature. *Prostaglandins* 51:225-232, 1996.
49. Nakayama T, Izumi Y, Soma M and Kanmatsuse K: Nitric oxide synthesis inhibitor prevents the ACTH-stimulated production of aldosterone in rat adrenal gland. *Endocrine Journal* 43:157-162, 1996.4
50. Kubo A, Fukuda N, Soma M, Izumi Y, and Kanmatsuse K: Inhibitory effect of an angiotensin II type 1 receptor antagonist on growth of vascular smooth muscle cells from spontaneously hypertensive rats. *J Cardiovascular Pharmacol* 27:58-63, 1996.

51. Nakayama T, Soma M, Izumi Y, and Kanmatsuse K: Organization of the human prostacyclin synthase gene. *Biochemical and Biophysical Research Communications* 221, 803-806, 1996.
52. Izumi Y, Matsuoka M, Kubo A, Nakayama T, Fukuda N, Watanabe Y, Soma M and Kanmatsuse K: Prevention of cardiac hypertrophy by a sub-antihypertensive dose of the α 1-adrenergic antagonist Bunazosin in Dahl salt-sensitive rats. *Hypertension Research* 19, 147-150, 1996.
53. 相馬正義、泉洋一、福田昇、渡辺吉康、上松瀬勝男：血圧日内変動からみた塩酸テモカプリル（エースコール）の降圧効果について。 *Geriat Med* 34:1695-1698, 1996.12.
54. Takahashi Y, Nakayama T, Soma M, Izumi Y and Kanmatsuse K: CA repeats polymorphism of the neuronal nitric oxide synthase gene. *Human Heredity* 47:58-59, 1997.
55. 相馬正義、福田昇、渡辺吉康、泉洋一、上松瀬勝男：血圧日内変動、腎機能、糖代謝、脂質代謝およびレニン-アンジオテンシン-アルドステロン系に対する塩酸イミダプリルの効果について。 *医学と薬学* 37:187-195, 1997.
56. Nakayama T, Soma M, Takahashi Y, Izumi Y, Kanmatsuse K and Esumi M: Association analysis of CA repeat polymorphism of endothelial nitric oxide synthase gene with essential hypertension in Japanese. *Clinical Genetics* 51:26-30, 1997.
57. 相馬正義、福田昇、泉洋一、渡辺吉康、上松瀬勝男：塩酸ベタキソロールの日内変動、レニン-アンジオテンシン-アルドステロン系、腎機能、糖代謝および脂質代謝に対する効果について。 *薬理と治療* 25: 1185-1193, 1997.
58. Nakayama T, Soma M, Takahashi Y, Uwabo J, Izumi Y, Kanmatsuse K. Novel polymorphic CA/TG repeat identified in the human prostacyclin synthase gene. *Human Heredity* 47: 176-177, 1997.
59. Takahashi Y, Nakayama T, Soma M, Uwabo J, Izumi Y, Kanmatsuse K. Association analysis of TG repeat polymorphism of the neuronal nitric oxide synthase gene with essential

- hypertension. *Clinical Genetics* 52:83-85, 1997.
60. Fukuda N, Soma M, Kubo A, Kunimoto M, Watanabe Y, Izumi Y and Kanmatsuse K. Effect of dibutyryl cyclic AMP on plasma renin activity in normal men and patients with primary aldosteronism. *J Endocrinol Invest* 20:13-17, 1997.
 61. Nakayama T, Soma M, Rahmudula D, Izumi Y, Kanmatsuse K. Nonsense mutation of prostacyclin synthase gene in a family. *Lancet* 349:1887-1888, 1997.6
 62. Fukuda N, Kubo A, Watanabe Y, Nakayama T, Soma M, Izumi Y, Kanmatsuse K. Antisense oligodeoxynucleotide complementary to platelet-derived growth factor A-chain messenger RNA inhibits the arterial proliferation in spontaneously hypertensive rats without altering their blood pressures. *J Hypertens* 15:1123-1136, 1997.10
 63. Fukuda N, Kishioka H, Satoh C, Nakayama T, Watanabe Y, Soma M, Izumi Y, Kanmatsuse K. Role of long-form PDGF A-chain in the growth of vascular smooth muscle cells from spontaneously hypertensive rats. *Am J Hypertens* 10:1117-1124, 1997.10
 64. Hu WY, Nakayama T, Fukuda N, Kishioka H, Soma M, Izumi Y, Kanmatsuse K. Sequences of exon 6 and the adjacent intron boundaries of the rat platelet-derived growth factor A-chain gene: Implications for alternative splicing. *Biochemical Genetics* 35:395-405, 1997.
 65. T. Nakayama, Izumi Y, Soma M, Kanmatsuse K. Adrenal renin-angiotensin-aldosterone system in streptozotocin-diabetic rats. *Horm Metab Res* 30:12-15, 1998.
 66. Uwabo J, Soma M, Nakayama T, Kanmatsuse K: Association of a variable number of tandem repeats in the endothelial constitutive nitric oxide synthase gene with essential hypertension in Japanese. *Am J Hypertens* 11:125-128, 1998.1
 67. 羽毛田 公、相馬正義、中山智祥、泉 洋一、上松瀬勝男：アンジオテンシノーゲン遺伝子多型 M235T と本態性高血圧症および左室肥大との関連性の検討。日大医学雑誌 57:12-16, 1998.1

68. 中山智祥、相馬正義、上松瀬勝男：ヒトプロスタサイクリン合成酵素遺伝子の5'上流領域の増幅と新しい多型の発見。日大医学雑誌 57:24-28, 1998.1
69. Fukuda N, Hu W.-Y, Kubo A, Endoh M, Kishioka H, Satoh T, Soma M, Izumi Y, Kanmatsuse K. Abnormal regulation of transforming growth factor- β receptor on vascular smooth muscle cells from spontaneously hypertensive rats by angiotensin II. *Hypertension* 31:672-677, 1998.4
70. Takahashi Y, Nakayama T, Soma M, Izumi Y, Kanmatsuse. Organization of the human natriuretic peptide receptor A gene. *Biochem Biophys Res Commun.* 246: 736-739, 1998.5
71. Kunitomo M. Soma M and Kanmatsuse K. Production of eicosanoid and angiotensin II in resistance vessels in spontaneously hypertensive rats. *Clin Experi Pharmacol Physiol* 25:430-434, 1998.7
72. Soma M, Nakayama T, Satoh M, Uwabo J, Duolikun R, Takahashi Y, Fukuda N, Watanabe Y, Izumi Y and Kanmatsuse K. A T1083C polymorphism in the human adenosine A2a receptor gene is not associated with essential hypertension. *Am J Hypertens* 11:1492-1494, 1998.12
73. 相馬正義、福田昇、泉洋一、渡辺吉康、上松瀬勝男：塩酸ブナゾシン徐放性製剤の降圧効果と心・腎・内分泌代謝系に及ぼす影響。医学と薬学、40:1047-1056, 1998.12
74. Hu W.-Y, Fukuda N, Soma M, Izumi Y, Kanmatsuse K. Bunazosine hydrochloride inhibits exaggerated growth of vascular smooth muscle cells from spontaneously hypertensive rats by suppressing the response to growth factors. *J Cardiovasc Pharmacol* 32:975-982, 1998.12
75. Duolikun R, Nakayama T, Soma M, Takahashi Y, Uwabo J, Sato M, Yoichi I, Kanmatsuse K, Ozawa Y. Structure of the type B human natriuretic peptide receptor gene and association of a novel microsatellite polymorphism with essential hypertension. *Cir Res* 84:605-610, 1999.
76. Fukuda N, Satoh T, Hu W.-Y, Soma M, Kubo A, Kishioka H, Watanabe Y, Izumi Y,

- Kanmatsuse K. Production of angiotensin II by homogenous cultures of vascular smooth muscle cells from spontaneously hypertensive rats. *Arterioscler Thromb Vascul Biol* 19:1210-1217, 1999.3
77. Nakayama M, Fukuda N, Watanabe Y, Soma M, Hu WY, Kishioka H, Satoh C, Kubo A, Kanmatsuse K. Low dose of eicosapentaenoic acid inhibits the exaggerated growth of vascular smooth muscle cells from spontaneously hypertensive rats through suppression of transforming growth factor-beta. *J Hypertens* 17:1421-30, 1999.10
78. Nakayama T, Soma M, Takahashi Y, Rehemudula D, Sato M, Uwabo J, Izumi Y, Kanmatsuse K. Nucleotide sequence of the 5'-flanking region of the type A human natriuretic peptide receptor gene and association analysis using a novel microsatellite in essential hypertension. *Am J Hypertens* 12:1144-1148, 1999.11
79. Fukuda N, Hu WY, Kubo A, Kishioka H, Satoh C, Soma M, Izumi Y, Kanmatsuse K. Angiotensin II upregulates transforming growth factor-beta type I receptor on rat vascular smooth muscle cells. *Am J Hypertens* 13(2):191-8, 2000.2
80. Rahmutula D, Nakayama T, Soma M, Takahashi Y, Uwabo J, Sato M, Izumi Y, Kanmatsuse K, Ozawa Y. An Insertion/Deletion Polymorphism in Intron 18 of the Type B Human Natriuretic Peptide Receptor Gene Is Not Associated with Cerebral Infarction. *Hypertens Res* 23(2): 173-176, 2000.3
81. Kawamura H, Jumabay M, Mitsubayasi H, Izumi Y, Soma M, Ozawa Y, Rehemudula D, Mahmut M, Hu Y, Aisa M, Cheng ZH, Wang Zhen. 24-hour blood pressure in Uygur, Kazakh and Han elderly subjects in China. *Hypertens Res* 23(2): 177-185, 2000.3
82. Nakayama T, Soma M, Takahashi Y, Rehemudula D, Kanmatsuse K, Furuya K. Functional deletion mutation of the 5'-flanking region of the type A human natriuretic peptide receptor gene and its association with essential hypertension and left ventricular hypertrophy in the Japanese. *Circ Res*. 2000 Apr 28;86(8):841-845.
83. 泉 洋一、福田 昇、中山智祥、渡辺吉康、相馬正義、上松瀬勝男。降圧薬 1 日飲

み忘れによる血圧の変化。新薬と臨床 49(6):601-606, 2000.6

84. Sato M, Soma M, Nakayama T, Kanmatsuse K. Dopamine D1 Receptor Gene Polymorphism is Associated with Essential Hypertension. *Hypertension* 36(2):183-186, 2000.8
85. Rahmutula D, Nakayama T, Soma M, Takahashi Y, Uwabo J, Sato M, Izumi Y, Saito S, Honye J, Kanmatsuse K, Ozawa Y. A C2077T polymorphism of the type B human natriuretic peptide receptor gene is not associated with myocardial infarction. *Med Sci Monit* 6(6): 1053-1055, 2000.12
86. Nakayama T, Soma M, Rehemudula D, Takahashi Y, Tobe H, Satoh M, Uwabo J, Kunimoto M, Kanmatsuse K. Association of 5' upstream promoter region of prostacyclin synthase gene variant with cerebral infarction. *Am J Hypertens* 13(12):1263-7, 2000.12
87. Nakayama T, Soma M, Takahashi Y, Rehemudula D, Tobe H, Sato M, Uwabo J, Kunimoto M, Izumi Y, Kanmatsuse K. Polymorphism of the promoter region of prostacyclin synthase gene is not related to essential hypertension. *Am J Hypertens*. 14(5 Pt 1):409-11, 2001.5
88. Nakayama T, Soma M, Rahmutula D, Ozawa Y, Kanmatsuse K. Isolation of the 5'-flanking region of genes by thermal asymmetric interlaced polymerase chain reaction. *Med Sci Monit*. 7(3):345-9, 2001.5
89. Rahmutula D, Nakayama T, Soma M, Sato M, Izumi Y, Kanmatsuse K, Ozawa Y. Systematic screening of type B human natriuretic peptide receptor gene polymorphisms and association with essential hypertension. *J Hum Hypertens* 15(7):471-4, 2001.7
90. Nakayama T, Soma M, Rahmutula D, Kanmatsuse K: Mutation of the human prostacyclin synthase gene. *Advances in Prostaglandin and Leukotriene Research* 65-68, 2001.
91. Nakayama T, Minato M, Nakagawa M, Soma M, Tobe H, Aoi N, Kosuge K, Sato M, Ozawa Y, Kanmatsuse K, Kokubun S: A novel mutation in the Ca²⁺-Sensing Receptor Gene in Familial Hypocalciuric Hypercalcemia. *Endocrine* 15: 277-282, 2001.

92. Soma M, Nakayama T, Rahmutula D, Uwabo J, Sato Mikano, Kunimoto M, Aoi N, Kosuge K and Kanmatsuse K. Ser9Gly Polymorphism in the Dopamine D3 Receptor Gene is not Associated with Essential Hypertension in the Japanese. *Med Sci Monit.* 8;CR1-4, 2002.1
93. Nakayama T, Soma M, Rahmutula D, Tobe H, Sato M, Uwabo J, Aoi N, Kosuge K, Kunimoto M, Kanmatsuse K, Kokubun S: Association study between a novel single nucleotide polymorphism of the promoter region of the prostacyclin synthase gene and essential hypertension. *Hypertens Res.* 2002 Jan;25(1):65-8.
94. Rahmutula D, Nakayama T, Soma M, Kosuge K, Aoi N, Izumi Yoichi, Kanmatsuse K and Ozawa Y: Structure and Polymorphisms of the Human Natriuretic Peptide Receptor C Gene. *Endocrine* 2002 17(2):85-90.
95. Nakayama T, Soma M, Saito S, Honye J, Yajima J, Rahmutula D, Kaneko Y, Sato M, Uwabo J, Aoi N, Kosuge K, Kunimoto M, Kanmatsuse K, Kokubun S. Association of a novel single nucleotide polymorphism of the prostacyclin synthase gene with myocardial infarction. *Am Heart J* 2002 May;143(5):797-801
96. Nakayama T, Soma M, Mizutani Y, Xinjuan Xu, Honye J, Kaneko Y, Rahmutula D, Aoi N, Kosuge K, Saito S, Kanmatsuse K, Kokubun S. A novel missense mutation of exon 3 in the type A human natriuretic peptide receptor gene: Possible association with essential hypertension. *Hypertens Res.* 2002 May;25(3):395-401.
97. Okada K, Onishi Y, Nagura Y, Soma M, Saito S, Kanmatsuse K, Takahashi S. Sclerosing encapsulating peritonitis: Regional changes of peritoneum. *Nephron* 2002 Oct;92(2):481-3
98. Nakayama T, Soma M, Watanabe Y, Hasimu B, Sato M, Aoi N, Kosuge K, Kanmatsuse K, Kokubun S, Marrow J, Oates J. Splicing mutation of the prostacyclin synthase gene in a family associated with hypertension. *Biochem Biophys Res Commun* 2002 Oct 11;297(5):1135-1139
99. 相馬正義、中山智祥、松本紘一、上松瀬勝男。塩酸バルニジピン長期投与の血圧日

内変動、レニン・アンジオテンシン・アルドステロン系、腎機能、糖代謝および脂質代謝に対する効果について。医学と薬学 49(1):93-101, 2003.1

100. Fukuda N, Nakayama M, Jian T, Satoh C, Nakayama T, Soma M, Izumi Y, Kanmatsuse K. Leukocyte angiotensin II levels in patients with essential hypertension: relation to insulin resistance. *Am J Hypertens* 2003 Feb;16(2):129-34.
101. Hasimu B, Nakayama T, Mizutani Y, Izumi Y, Asai S, Soma M, Kokubun S, Ozawa Y. Haplotype analysis of the human renin gene and essential hypertension. *Hypertension* 2003 Feb;41(2):308-12
102. Nakayama T, Soma M, Watanabe Y, Hasimu B, Kanmatsuse K, Kokubun S, Morrow JD, Oates JA. Splicing mutation of the prostacyclin synthase gene in a family associated with hypertension. *Adv Exp Med Biol* 2003;525:165-8
103. Hasimu B, Nakayama T, Mizutani Y, Izumi Y, Asai S, Soma M, Kokubun S, Ozawa Y. A novel variable number of tandem repeat polymorphism of the renin gene and essential hypertension. *Hypertens Res.* 2003 Jun;26(6):473-7.
104. Takahiro Ueno, Emi Shimazaki, Taro Matsumoto, Hideyuki Watanabe, Akiko Tsunemi, Yumiko Takahashi, Mayumi Mori, Rumi Hamano, Toshiyuki Fujioka, Masayoshi Soma, Koichi Matsumoto, Katsuo Kanmatsuse. Paraoxonase1 polymorphism Leu-Met55 is associated with cerebral infarction in Japanese population. *Med Sci Monit*, 2003; 9(6): CR260-264
105. Nakayama T, Soma M, Saito S, Honye J, Sato M, Aoi N, Kosuge K, Haketa A, Kanmatsuse K, Kokubun S. Missense mutation of exon 3 in the type A human natriuretic peptide receptor gene is associated with myocardial infarction. *Med Sci Monit*, 2003, 9(12):CR505-510
106. Nakayama T, Soma M, Haketa A, Aoi N, Kosuge K, Sato M, Kanmatsuse K, Kokubun S. Haplotype analysis of the prostacyclin synthase gene and essential hypertension. *Hypertens Res.* 2003 Jul;26(7):553-7

107. Nakayama T, Soma M, Watanabe Y, Hasimu B, Kanmatsuse K, Kokubun S, Morrow JD, Oates JA. Splicing mutation of the prostacyclin synthase gene in a family associated with hypertension. *Adv Exp Med Biol.* 2003;525:165-8.
108. Nakayama T, Soma M, Kanmatsuse K, Kokubun S. The microsatellite alleles on chromosome 1 associated with essential hypertension and blood pressure levels. *J Hum Hypertens.* 2004 Nov;18(11):823-8.
109. Haketa A, Soma M, Nakayama T, Sato M, Kosuge K, Aoi N, Matsumoto K. Two medium-chain acyl-coenzyme A synthetase genes, SAH and MACS1, are associated with plasma high-density lipoprotein cholesterol levels, but they are not associated with essential hypertension. *J Hypertens.* 2004 Oct;22(10):1903-7.
110. Nakayama T, Soma M, Sato N, Haketa A, Kosuge K, Aoi N, Sato M, Izumi Y, Matsumoto K, Kanmatsuse K, Kokubun S. An association study in essential hypertension using functional polymorphisms in lymphotoxin-alpha gene. *Am J Hypertens.* 2004 Nov;17(11 Pt 1):1045-9.
111. Higuchi T, Yamamoto C, Kuno T, Okada K, Soma M, Fukuda N, Nagura Y, Takahashi S, Matsumoto K. A comparison of bicarbonate hemodialysis hemodiafiltration, and acetate-free biofiltration on cytokine production. *Therapeutic Apheresis and Dialysis* 8(6):460-467, 2004.
112. 徳竹英一、相馬正義 :HMG-CoA 還元酵素阻害薬と長時間作用型 Ca 拮抗薬との併用による高脂血症を合併した高血圧患者の頸動脈内膜中膜肥厚度(IMT)の改善効果。 *Therapeutic Res* 26(1):145-152, 2005.
113. Aoi N, Soma M, Nakayama T, Rahmutula D, Kosuge K, Izumi Y, Matsumoto K. Variable number of tandem repeat of the 5'-flanking region of type-C human natriuretic peptide receptor gene influences blood pressure levels in obesity-associated hypertension. *Hypertens Res.* 2004 Oct;27(10):711-6.

114. Sano M, Kuroi N, Nakayama T, Sato N, Izumi Y, Soma M, Kokubun S. Association study of calcitonin-receptor-like receptor gene in essential hypertension. *Am J Hypertens*. 2005 Mar;18(3):403-8.
115. Kushiro T, Takahashi A, Saito F, Otsuka Y, Soma M, Kurihara T, Satomura A, Saito T, Kanmatsuse K. Erectile dysfunction and its influence on quality of life in patients with essential hypertension. *Am J Hypertens*. 2005 Mar;18(3):427-30.
116. Kobayashi Y, Nakayama T, Sato N, Izumi Y, Kokubun S, Soma M. Haplotype-based case-control study revealing an association between the adrenomedullin gene and proteinuria in subjects with essential hypertension. *Hypertens Res*. 2005 Mar;28(3):229-36.
117. Sano M, Kuroi N, Nakayama T, Sato N, Izumi Y, Soma M. Association study: the aminopeptidase A gene and essential hypertension. *International Journal of Biomedical Science (IJBS)*. 1 :16-22.
118. Nakayama T, Asai S, Sato N, Soma M. Genotype and Haplotype Association Study of the STRK1 Region on 5q12 Among Japanese. A Case-Control Study. *Stroke* 2006 Jan;37(1):69-76
119. Sato N, Nakayama T, Asai S, Soma M. A haplotype in the human Sur2 gene is associated with essential hypertension. *J Human Hypertension* 2006, 20(1):87-90
120. Higuchi T, Fukuda N, Yamamoto C, Yamazaki T, Oikawa O, Ohnishi Y, Soma M, Matsumoto K. The influence of uremic serum on interleukin-1 β and interleukin-1 receptor antagonist production by peripheral blood mononuclear cells. *Therapeutic Apheresis and Dialysis* 10(1):65-71, 2006
121. Higuchi T, Matsukawa Y, Okada K, Oikawa O, Yamazaki T, Ohnishi Y, Fujita T, Fukuda N, Soma M, Matsumoto K. Correction of Copper Deficiency Improves Erythropoietin Unresponsiveness in Hemodialysis Patients with Anemia. *Intern Med* 2006 May;45(5):271-3

122. Sato M, Soma M, Nakayama T, Kosuge K, Suzuki R, Okada K, Komatsu K, Sugitani M, Matsumoto K. A case of adrenocorticotropin-independent bilateral adrenal macronodular hyperplasia (AIMAH) with primary hyperparathyroidism (PHPT). *Endocrine Journal*. 53: 111-117
123. Morita A, Nakayama T, Soma M. Association study between C-reactive protein genes and ischemic stroke in Japanese subjects. *Am J Hypertens* 2006 Jun;19(6):593-600.
124. Morita A, Nakayama T, Doba N, Hinohara S, Soma M. Polymorphism of the C-reactive protein (CRP) gene is related to serum CRP Level and arterial pulse wave velocity in healthy elderly Japanese. *Hypertens Res*. 2006 May ; 29(5):323-31.
125. Nakayama T, Kuroi N, Sano M, Tabara Y, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T, Soma M. Mutation of the follicle-stimulating hormone receptor gene 5'-untranslated region associated with female hypertension. *Hypertension*. 2006 Sep;48(3):512-8. Epub 2006 Jul 24.
126. Omi T, Kumada M, Kamesaki T, Okuda H, Munkhtulga L, Yanagisawa Y, Utsumi N, Gotoh T, Hata A, Soma M, Umemura S, Ogihara T, Takahashi N, Tabara Y, Shimada K, Mano H, Kajii E, Miki T, Iwamoto S. An intronic variable number of tandem repeat polymorphisms of the cold-induced autoinflammatory syndrome 1 (CIAS1) gene modifies gene expression and is associated with essential hypertension. *Eur J Hum Genet*. 2006 Jul 26; [Epub ahead of print]
127. Saito K, Nakayama T, Sato N, Morita A, Takahashi T, Soma M, Usami R. Haplotypes of the plasminogen activator gene associated with ischemic stroke. *Thromb Haemost* 2006 Sep;96(3):331-6.
128. Morita A, Nakayama T, Doba N, Hinohara S, Mizutani T, Soma M. Genotyping of triallelic SNPs using TaqMan((R)) PCR. *Mol Cell Probes*. 2006 Nov 7.
129. 小牧宏一、相馬正義、坂田悍教、小菅琴子、青井則子、羽毛田公、中山智祥、岡本順

子、細川武、土居通哉、五味敏明、松本絃一：高齢者住民検診における血漿高分子量
アディポネクチン濃度。埼玉県立大学 紀要 8:33-37, 2006

130. Kaneko Y, Nakayama T, Saito K, Morita A, Sato I, Maruyama A, Soma M, Takahashi T, Sato N. Relationship between the thromboxane A2 receptor gene and susceptibility to cerebral infarction. *Hypertens Res.* 2006 Sep;29(9):665-71.
131. Saito K, Nakayama T, Sato N, Kaneko Y, Sato I, Morita A, Takahashi T, Soma M, Usami R. Haplotype based association study between t-PA gene and essential hypertension. *International Journal of Biomedical Science (IJBS)*. 2: 166-171
132. Nakayama T, Asai S, Sato N, Soma M. PDE4D gene in the STRK1 region on 5q12: susceptibility gene for ischemic stroke. *Curr Med Chem.* 2007;14(30):3171-8.
133. Higuchi T, Yamazaki T, Ohnishi Y, Matsumoto S, Yabuki M, Kitajima S, Toshida Y, Maruyama N, Oikawa O, Okada K, Fukuda N, Soma M, Matsumoto. A case report of hemodialysis intolerance with eosinophilia. *Ther Apher Dial.* 2007 Feb;11(1):70-3.
134. Yatsu K, Mizuki N, Hirawa N, Oka A, Itoh N, Yamane T, Ogawa M, Shiwa T, Tabara Y, Ohno S, Soma M, Hata A, Nakao K, Ueshima H, Ogihara T, Tomoike H, Miki T, Kimura A, Mano S, Kulski JK, Umemura S, Inoko H. High-resolution mapping for essential hypertension using microsatellite markers. *Hypertension.* 2007 Mar;49(3):446-52.
135. Aoi N, Nakayama T, Tahira Y, Haketa A, Yabuki M, Sekiyama T, Nakane C, Mano H, Kawachi H, Sato N, Soma M, Matsumoto K. Two novel genotypes of the thiazide-sensitive Na-Cl cotransporter (SLC12A3) gene in patients with Gitelman's syndrome. *Endocrine.* 2007 Apr;31(2):149-53.
136. Ueno T, Takahashi Y, Matsumoto T, Tsunemi A, Watanabe H, Tahira K, Fukuda N, Soma M, Matsumoto K. Postprandial plasma lipid levels are influenced by the interaction of functional polymorphisms in the microsomal triglyceride transfer protein and β 3 adrenergic receptor genes. *Med Sci Monit.* 2007; 13(5): BR112-118

137. Morita A, Nakayama T, Soma M, Mizutani T. Association Between the Calcitonin-Related Peptide alpha (CALCA) Gene and Essential Hypertension in Japanese. *Am J Hypertens*. 2007 May; 20(5):527-32.
138. Kosuge K, Soma M, Nakayama T, Aoi N, Sato M, Izumi Y, Matsumoto K. A novel variable number of tandem repeat of the natriuretic peptide precursor B gene's 5'-flanking region is associated with essential hypertension among Japanese females. *International Journal of Medical Sciences*. 2007 May 16;4(3):146-52.
139. Sato M, Nakayama T, Soma M, Aoi N, Kosuge K, Haketa A, Izumi Y, Matsumoto K, Sato N, Kokubun S. Association between prostaglandin E2 receptor gene and essential hypertension. *Prostaglandins Leukot Essent Fatty Acids*. 2007 Jul;77(1):15-20.
140. Hui P, Nakayama T, Morita A, Sato N, Hishiki M, Saito K, Yoshikawa Y, Tamura M, Sato I, Takahashi T, Soma M, Izumi Y, Ozawa Y, and Cheng Z. Common Single Nucleotide Polymorphisms in Japanese Patients with Essential Hypertension: Aldehyde Dehydrogenase 2 Gene as a Risk Factor Independent of Alcohol Consumption. *Hypertens Res* 2007 Jul; 30(7): 585-592
141. Yoshikawa Y, Nakayama T, Saito K, Hui P, Morita A, Sato N, Takahashi T, Tamura M, Sato I, Aoi N, Doba N, Hinohara S, Soma M, Usami R. Haplotype-based case-control study of the association between the guanylate cyclase activator 2B (GUCA2B, Uroguanylin) gene and essential hypertension. *Hypertens Res*. 2007 Sep;30(9):789-96.
142. Shimodaira M, Nakayama T, Sato N, Saito K, Morita A, Sato I, Takahashi T, Soma M, Izumi Y. Association study of aromatase gene (CYP19A1) in essential hypertension. *Int J Med Sci*. 2008 Feb 7;5(1):29-35.
143. Kato N, Miyata T, Tabara Y, Katsuya T, Yanai K, Hanada H, Kamide K, Nakura J, Kohara K, Takeuchi F, Mano H, Yasunami M, Kimura A, Kita Y, Ueshima H, Nakayama T, Soma M, Hata A, Fujioka A, Kawano Y, Nakao K, Sekine A, Yoshida T, Nakamura Y, Saruta T, Ogihara T, Sugano S, Miki T, Tomoike H. High-density association study and nomination of susceptibility genes for hypertension in the Japanese National Project. *Hum Mol Genet*.

2008 Feb 15;17(4):617-27.

144. Kohara K, Tabara Y, Nakura J, Imai Y, Ohkubo T, Hata A, Soma M, Nakayama T, Umemura S, Hirawa N, Ueshima H, Kita Y, Ogihara T, Katsuya T, Takahashi N, Tokunaga K, and Miki T. Identification of Hypertension-Susceptibility Genes and Pathways by a Systemic Multiple Candidate Gene Approach: The Millennium Genome Project for Hypertension. *Hypertens Res*. 2008 Feb;31(2):203-12.
145. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Ozawa Y, Ma Y. A haplotype of the CYP4A11 gene associated with essential hypertension in Japanese men. *J Hypertens*. 2008 Mar; 26(3):453-61.
146. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Matsumoto K, Ozawa Y, Ma Y. Haplotype-based case study of human CYP4A11 gene and cerebral infarction in Japanese subject. *Endocrine*. 2008 Apr; 33(2):215-22.
147. Mashimo Y, Suzuki Y, Hatori K, Tabara Y, Miki T, Tokunaga K, Katsuya T, Ogihara T, Yamada M, Takahashi N, Maikita Y, Nakayama T, Soma M, Hirawa N, Umemura S, Ohkubo T, Imai Y and Hata A. Association of TNFRSF4 gene polymorphisms with essential hypertension. *Journal of Hypertension*. 2008 May;26(5):902-13.
148. Aoi N, Soma M, Nakayama T, Kosuge K, Haketa A, Sato M, Sato N. Case-control study of the role of the Gitelman's syndrome gene in essential hypertension. *Endocrine Journal*. 2008 May;55(2):305-10.
149. Ueno T, Soma M, Tabara Y, Tokunaga K, Tahira K, Fukuda N, Matsumoto K, Nakayama T, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T. Association between fatty acid binding protein 3 gene variants and essential hypertension in humans. *Am J Hypertens*. 2008 Jun; 21(6):691-5.
150. Naganuma T, Nakayama T, Sato N, Fu Z, Soma M, Aoi N, Usami R. A haplotype-based case-control study examining human extracellular superoxide dismutase gene and essential hypertension. *Hypertens Res*. 2008 Aug; 31(8):1533-40.

151. Kosuge K, Soma M, Nakayama T, Aoi N, Sato M, Haketa A, Uwabo J, Izumi Y, Matsumoto K. Human Uncoupling Protein 2 and 3 Genes Are Associated with Obesity in Japanese. *Endocrine*. 2008 Dec;34(1-3):87-95.
152. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Matsumoto K, Ozawa Y, Ma Y. Haplotype-based case-control study of the human CYP4F2 gene and essential hypertension in Japanese subjects. *Hypertens Res*. 2008 Sep; 31(9):1719-26.
153. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Matsumoto K, Ozawa Y, Ma Y. A haplotype of the CYP4F2 gene is associated with cerebral infarction in Japanese men. *Am J Hypertens*. 2008 Nov; 21(11):1216-23.
154. Naganuma T, Nakayama T, Sato N, Fu Z, Soma M, Aoi N, Hinohara S, Doba N, Usami R. Association of extracellular superoxide dismutase gene with cerebral infarction in women: a haplotype-based case-control study. *Hereditas*. 2008 Dec;145(6):283-92.
155. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Ozawa Y, Ma Y, Matsumoto K, Doba N, Hinohara S. A haplotype of the CYP4F2 gene associated with myocardial infarction in Japanese men. *Mol Genet Metab*. 2009 Mar;96(3):145-7.
156. Ueno T, Fukuda N, Nagase H, Tsunemi A, Tahira K, Matsumoto T, Hiraoka-Yamamoto J, Ikeda K, Mitsumata M, Sato Y, Soma M, Matsumoto K, Yamori Y. Atherogenic dyslipidemia and altered hepatic gene expression in SHRSP.Z-Leprfa/IzmDmcr rats. *Int J Mol Med*. 2009 Mar;23(3):313-20.
157. Ueno T, Tabara Y, Fukuda N, Tahira K, Matsumoto T, Kosuge K, Haketa A, Matsumoto K, Sato Y, Nakayama T, Katsuya T, Ogihara T, Makita Y, Hata A, Yamada M, Takahashi N, Hirawa N, Umemura S, Miki T, Soma M. Association of SLC6A9 gene variants with human essential hypertension. *J Atheroscler Thromb*. 2009 Jun;16(3):201-6.

158. Ueno T, Watanabe H, Fukuda N, Tsunemi A, Tahira K, Matsumoto T, Takayama T, Chiku M, Saito S, Sato Y, Hirayama A, Matsumoto K, Soma M. Influence of genetic polymorphisms in oxidative stress related genes and smoking on plasma MDA-LDL, soluble CD40 ligand, E-selectin and soluble ICAM1 levels in patients with coronary artery disease. *Med Sci Monit.* 2009 Jul;15(7):CR341-8.
159. Yamaguchi M, Nakayama T, Fu Z, Naganuma T, Sato N, Soma M, Doba N, Hinohara S, Morita A, Mizutani T. Relationship between haplotypes of KCNN4 gene and susceptibility to human vascular diseases in Japanese. *Med Sci Monit.* 2009 Aug;15(8):CR389-97.
160. Naganuma T, Nakayama T, Sato N, Fu Z, Yamaguchi M, Soma M, Aoi N, Usami R, Doba N, Hinohara S. Haplotype-based case-control study between human apurinic/aprimidinic endonuclease 1/redox effector factor-1 gene and cerebral infarction. *Clin Biochem.* 2009 Oct;42(15):1493-9.
161. Wang Z, Nakayama T, Sato N, Yamaguchi M, Izumi Y, Kasamaki Y, Ohta M, Soma M, Aoi N, Ozawa Y, Ma Y, Doba N, Hinohara S. Purinergic receptor P2Y, G-protein coupled, 2 (P2RY2) gene is associated with cerebral infarction in Japanese subjects. *Hypertens Res.* 2009 Nov;32(11):989-96.
162. Nakayama T, Aoi N, Sato N, Sato M, Kosuge K, Izumi Y, Soma M, Matsumoto K. [Establishment of genetic testing for Gitelman's syndrome]. *Rinsho Byori.* 2010 Feb;58(2):156-61.
163. Wang ZX, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Ohta M, Soma M, Aoi N, Matsumoto K, Ozawa Y, Ma YT, Doba N, Hinohara S. Association of the Purinergic Receptor P2Y, G-Protein Coupled, 2 (P2RY2) Gene With Myocardial Infarction in Japanese Men. *Circ J.* 2009 Dec;73(12):2322-9.
164. Naganuma T, Nakayama T, Sato N, Fu Z, Soma M, Yamaguchi M, Shimodaira M, Aoi N, Usami R. Haplotype-based case-control study on human apurinic/aprimidinic

endonuclease 1/redox effector factor-1 gene and essential hypertension. *Am J Hypertens.* 2010 Feb;23(2):186-91.

165. Wang Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Ohta M, Soma M, Aoi N, Ozawa Y, Ma Y. The purinergic receptor P2Y, G-protein coupled, 2 (P2RY2) gene associated with essential hypertension in Japanese men. *J Hum Hypertens.* 2010 May;24(5):327-35.
166. Nakazato T, Nakayama T, Naganuma T, Sato N, Fu Z, Wang Z, Soma M, Sugama K, Hinohara S, Doba N. Haplotype-based case-control study of receptor (calcitonin) activity-modifying protein-1 gene in cerebral infarction. *J Hum Hypertens.* 2010 May;24(5):351-8.
167. Shimodaira M, Nakayama T, Sato N, Naganuma T, Yamaguchi M, Aoi N, Sato M, Izumi Y, Soma M, Matsumoto K. Association study of the elastin microfibril interfacier 1 (EMILIN1) gene in essential hypertension. *Am J Hypertens.* 2010 May;23(5):547-55.
168. Abe M, Okada K, Maruyama T, Maruyama N, Soma M, Matsumoto K. Clinical effectiveness and safety evaluation of long-term pioglitazone treatment for erythropoietin responsiveness and insulin resistance in type 2 diabetic patients on hemodialysis. *Expert Opin Pharmacother.* 2010 Jul;11(10):1611-20.
169. Nishida Y, Takahashi Y, Nakayama T, Soma M, Kitamura N, Asai S. Effect of candesartan monotherapy on lipid metabolism in patients with hypertension: a retrospective longitudinal survey using data from electronic medical records. *Cardiovasc Diabetol.* 2010 Aug 16;9:38.
170. Abe M, Okada K, Maruyama N, Matsumoto S, Maruyama T, Fujita T, Matsumoto K, Soma M. Comparison between the antiproteinuric effects of the calcium channel blockers benidipine and cilnidipine in combination with angiotensin receptor blockers in hypertensive patients with chronic kidney disease. *Expert Opin Investig Drugs.* 2010 Sep;19(9):1027-37.
171. Aoi N, Nakayama T, Soma M, Kosuge K, Haketa A, Sato M, Sato N, Asai S, Matsumoto K.

Association of the insulin-like growth factor1 gene with myocardial infarction in Japanese subjects. *Hereditas*. 2010 Oct;147(5):215-24.

172. Shimodaira M, Nakayama T, Sato N, Aoi N, Sato M, Izumi Y, Soma M, Matsumoto K. Association of HSD3B1 and HSD3B2 gene polymorphisms with essential hypertension, aldosterone level, and left ventricular structure. *Eur J Endocrinol*. 2010 Oct;163(4):671-80.
173. Tabara Y, Kohara K, Kita Y, Hirawa N, Katsuya T, Ohkubo T, Hiura Y, Tajima A, Morisaki T, Miyata T, Nakayama T, Takashima N, Nakura J, Kawamoto R, Takahashi N, Hata A, Soma M, Imai Y, Kokubo Y, Okamura T, Tomoike H, Iwai N, Ogihara T, Inoue I, Tokunaga K, Johnson T, Caulfield M, Munroe P; Global Blood Pressure Genetics Consortium, Umemura S, Ueshima H, Miki T. Common variants in the ATP2B1 gene are associated with susceptibility to hypertension: the Japanese Millennium Genome Project. *Hypertension*. 2010 Nov;56(5):973-80.
174. Yamaguchi M, Nakayama T, Fu Z, Sato N, Soma M, Morita A, Hinohara S, Doba N, Mizutani T. The haplotype of the CACNA1B gene associated with cerebral infarction in a Japanese population. *Hereditas*. 2010 Dec;147(6):313-9.
175. Yoshida Y, Fukuda N, Maeshima A, Yamamoto C, Matsumoto T, Ueno T, Nojima Y, Matsumoto K, Soma M. Treatment with valsartan stimulates endothelial progenitor cells and renal label-retaining cells in hypertensive rats. *J Hypertens*. 2011 Jan;29(1):91-101.
176. Abe M, Maruyama N, Okada K, Matsumoto S, Matsumoto K, Soma M. Effects of Lipid-Lowering Therapy with Rosuvastatin on Kidney Function and Oxidative Stress in Patients with Diabetic Nephropathy. *J Atheroscler Thromb*. 2011;18(11):1018-28.
177. Abe M, Maruyama N, Matsumoto S, Okada K, Fujita T, Matsumoto K, Soma M. Comparison of Sustained Hemodiafiltration with Acetate-Free Dialysate and Continuous Venovenous Hemodiafiltration for the Treatment of Critically Ill Patients with Acute Kidney Injury. *Int J Nephrol*. 2011

178. Abe M, Okada K, Maruyama T, Maruyama N, Matsumoto K, Soma M. Relationship between erythropoietin responsiveness, insulin resistance, and malnutrition-inflammation-atherosclerosis (MIA) syndrome in hemodialysis patients with diabetes. *Int J Artif Organs*. 2011 Jan;34(1):16-25.
179. Abe M, Maruyama N, Yoshida Y, Ito M, Okada K, Soma M. Efficacy analysis of the lipid-lowering and renoprotective effects of rosuvastatin in patients with chronic kidney disease. *Endocr J*. 2011;58(8):663-74.
180. Ito M, Abe M, Okada K, Sasaki H, Maruyama N, Tsuchida M, Higuchi T, Kikuchi F, Soma M. The dipeptidyl peptidase-4 (DPP-4) inhibitor vildagliptin improves glycemic control in type 2 diabetic patients undergoing hemodialysis. *Endocr J*. 2011;58(11):979-87.
181. Abe M, Okada K, Soma M. Antidiabetic agents in patients with chronic kidney disease and end-stage renal disease on dialysis: metabolism and clinical practice. *Curr Drug Metab*. 2011 Jan;12(1):57-69.
182. Abe M, Okada K, Soma M, Matsumoto K. Relationship between insulin resistance and erythropoietin responsiveness in hemodialysis patients. *Clin Nephrol*. 2011 Jan;75(1):49-58.
183. Abe M, Okada K, Maruyama N, Matsumoto S, Maruyama T, Fujita T, Matsumoto K, Soma M. Benidipine reduces albuminuria and plasma aldosterone in mild-to-moderate stage chronic kidney disease with albuminuria. *Hypertens Res*. 2011 Feb;34(2):268-73.
184. Yamamoto C, Fukuda N, Jumabay M, Saito K, Matsumoto T, Ueno T, Soma M, Matsumoto K, Shimosawa T. Protective effects of statin on cardiac fibrosis and apoptosis in adrenomedullin-knockout mice treated with angiotensin II and high salt loading. *Hypertens Res*. 2011 Mar;34(3):348-53.
185. Abe M, Okada K, Ikeda K, Matsumoto S, Soma M, Matsumoto K. Characterization of

- insulin adsorption behavior of dialyzer membranes used in hemodialysis. *Artif Organs*. 2011 Apr;35(4):398-403.
186. Fujita T, Shimizu C, Fuke Y, Satomura A, Abe M, Kaizu K, Matsumoto K, Soma M. Serum interleukin-18 binding protein increases with behavior different from IL-18 in patients with diabetic nephropathy. *Diabetes Res Clin Pract*. 2011 Jun;92(3):e66-9.
187. Sezai A, Soma M, Hata M, Yoshitake I, Unosawa S, Wakui S, Shiono M. Effects of Olmesartan on the Renin-angiotensin-aldosterone System for Patients with Essential Hypertension after Cardiac Surgery-Investigation Using a Candesartan Change-over Study. *Ann Thorac Cardiovasc Surg*. 2011;17(5):487-93.
188. Matsumoto K, Shimodaira M, Nakagawa T, Nakayama T, Nakazato T, Izumi Y, Soma M, Matsumoto K, Sato N, Aoi N. Association study: SLC6A18 gene and myocardial infarction. *Clin Biochem*. 2011 Jul;44(10-11):789-94.
189. Abe M, Maruyama N, Okada K, Matsumoto S, Matsumoto K, Soma M. Additive antioxidative effects of azelnidipine on angiotensin receptor blocker olmesartan treatment for type 2 diabetic patients with albuminuria. *Hypertens Res*. 2011 Aug;34(8):935-41.
190. 藤田之彦, 橋本 修, 相馬正義, 竹内 仁, 相澤 信, 片山容一. 学生の講義出席率からみたハイブリッド型 PBL tutorial 導入の効果. *日大医学雑誌*. 2011 Oct;70(5):244-248.
191. Yamazaki D, Tabara Y, Kita S, Hanada H, Komazaki S, Naitou D, Mishima A, Nishi M, Yamamura H, Yamamoto S, Kakizawa S, Miyachi H, Yamamoto S, Miyata T, Kawano Y, Kamide K, Ogihara T, Hata A, Umemura S, Soma M, Takahashi N, Imaizumi Y, Miki T, Iwamoto T, Takeshima H. TRIC-A channels in vascular smooth muscle contribute to blood pressure maintenance. *Cell Metab*. 2011 Aug 3;14(2):231-41.

192. Nishida Y, Takahashi Y, Nakayama T, Soma M, Asai S. Comparative effect of olmesartan and candesartan on lipid metabolism and renal function in patients with hypertension: a retrospective observational study. *Cardiovasc Diabetol*. 2011 Aug 10;10:74.
193. Abe M, Maruyama T, Fujii Y, Kitai M, Okada K, Matsumoto K, Soma M. Disopyramide-induced hypoglycemia in a non-diabetic hemodialysis patient: a case report and review of the literature. *Clin Nephrol*. 2011 Nov;76(5):401-6.
194. Tahira K, Ueno T, Fukuda N, Aoyama T, Tsunemi A, Matsumoto S, Nagura C, Matsumoto T, Soma M, Shimba S, Matsumoto Y. Obesity alters the expression profile of clock genes in peripheral blood mononuclear cells. *Arch Med Sci*. 2011 Dec 31;7(6):933-40.
195. Kajiwara M, Ueno T, Fukuda N, Matsuda H, Shimokawa T, Kitai M, Tsunemi A, Fuke Y, Fujita T, Matsumoto K, Matsumoto Y, Ra C, Soma M. Development of pyrrole-imidazole polyamide targeting fc receptor common gamma chain for the treatment of immune-complex related renal disease. *Biol Pharm Bull*. 2012;35(11):2028-35.
196. Shimizu C, Fujita T, Fuke Y, Ito K, Satomura A, Matsumoto K, Soma M. High circulating levels of interleukin-18 binding protein indicate the severity of glomerular involvement in systemic lupus erythematosus. *Mod Rheumatol*. 2012 Feb;22(1):73-9.
197. Han Y, Fukuda N, Ueno T, Endo M, Ikeda K, Xueli Z, Matsumoto T, Soma M, Matsumoto K. Role of Complement 3a in the Synthetic Phenotype and Angiotensin II-Production in Vascular Smooth Muscle Cells From Spontaneously Hypertensive Rats. *Am J Hypertens*. 2012 Mar;25(3):284-9.
198. Tabara Y, Kohara K, Miki T; Millennium Genome Project for Hypertension. Fujioka A, Hanada H, Hata A, Hirawa N, Hiura Y, Imai Y, Inoko H, Itoh N, Iwai N, Kulski JK, Kamide K, Kato N, Osaka TK, Kawamoto R, Kawano Y, Kimura A, Kita Y, Kohara K, Kokubo Y, Mano H, Mano S, Miki T, Miyata T, Mizuki N, Morisaki T, Nakamura Y, Nakao K, Nakayama T, Nakura

- J, Ogawa M, Ogihara T, Ohkubo T, Ohno S, Oka A, Okamura T, Saruta T, Sekine A, Shiwa T, Soma M, Sugano S, Tabara Y, Tajima A, Takahashi N, Takashima N, Takeuchi F, Tokunaga K, Tomoike H, Umemura S, Yamane T, Yanai K, Yasunami M, Yatsu K, Yoshida T, Tabara Y. Hunting for genes for hypertension: the Millennium Genome Project for Hypertension. *Hypertens Res.* 2012 Jun;35(6):567-73.
199. Fu Z, Nakayama T, Sato N, Izumi Y, Kasamaki Y, Shindo A, Ohta M, Soma M, Aoi N, Sato M, Ozawa Y, Ma Y. Haplotype-based case-control study of CYP4A11 gene and myocardial infarction. *Hereditas.* 2012 Jun;149(3):91-98.
200. Abe M, Maruyama N, Suzuki H, Fujii Y, Ito M, Yoshida Y, Okada K, Soma M. Additive renoprotective effects of aliskiren on angiotensin receptor blocker and calcium channel blocker treatments for type 2 diabetic patients with albuminuria. *Hypertens Res.* 2012 Aug;35(8):874-81.
201. 内田俊也, 兒島憲一郎, 種本雅之, 菅原正弘, 高橋正人, 齋藤友昭, 伊藤孝史, 相馬正義, J-Circle 研究会. N型Ca拮抗薬シルニジピンが尿酸産生に及ぼす影響について. *痛風と核酸代謝.* 2012;36(1):67
202. Jiang J, Nakayama T, Shimodaira M, Sato N, Aoi N, Sato M, Izumi Y, Kasamaki Y, Ohta M, Soma M, Matsumoto K, Kawamura H, Ozawa Y, Ma Y. A haplotype of the SMTN gene associated with myocardial infarction in Japanese women. *Genet Test Mol Biomarkers.* 2012 Sep;16(9):1019-26.
203. Jiang J, Nakayama T, Shimodaira M, Sato N, Aoi N, Sato M, Izumi Y, Kasamaki Y, Ohta M, Soma M, Matsumoto K, Kawamura H, Ozawa Y, Hinohara S, Doba N, Ma Y. Association of the smoothelin (SMTN) gene with cerebral infarction in men: a haplotype-based case-control study. *Vasc Med.* 2012 Oct;17(5):317-25.
204. Aoi N, Nakayama T, Soma M, Kosuge K, Haketa A, Sato M, Sato N, Hinohara S, Doba N,

- Asai S. The insulin-like growth factor-1 gene is associated with cerebral infarction in Japanese subjects. *Hereditas*. 2012 Oct;149(5):153-162.
205. Jiang J, Nakayama T, Shimodaira M, Sato N, Aoi N, Sato M, Izumi Y, Kasamaki Y, Ohta M, Soma M, Matsumoto K, Kawamura H, Ozawa Y, Ma Y. Haplotype of smoothelin gene associated with essential hypertension. *Hereditas*. 2012 Oct;149(5):178-185.
206. Abe M, Maruyama N, Suzuki H, Okada K, Soma M. International normalized ratio decreases after hemodialysis treatment in patients treated with warfarin. *J Cardiovasc Pharmacol*. 2012 Dec;60(6):502-7.
207. 藤井由季, 阿部雅紀, 樋口輝美, 鈴木紘子, 松本史郎, 水野真理, 菊池 史, 岡田一義, 相馬正義. 糖尿病血液透析患者におけるアログリプチンの有効性と安全性の検討. *透析会誌* 46(7):633-640.
208. Haketa A, Soma M, Nakayama T, Kosuge K, Aoi N, Hishiki M, Hatanaka Y, Ueno T, Doba N, Hinohara S. Association between SIRT2 gene polymorphism and height in healthy, elderly Japanese subjects. *Transl Res*. 2013 Jan;161(1):57-8.
209. Sezai A, Soma M, Nakata KI, Hata M, Yoshitake I, Wakui S, Hata H, Shiono M. Comparison of Febuxostat and Allopurinol for Hyperuricemia in Cardiac Surgery Patients (NU-FLASH Trial). *Circ J*. 2013;77(8):2043-9.
210. Kato K, Ishii Y, Mazaki T, Uehara T, Nakamura H, Kikuchi H, Yamagami H, Sato H, Mizuno S, Soma M, Henmi A, Masuda H, Moriyama M, Tanaka M. Spontaneous Regression of Polyposis following Abdominal Colectomy and Helicobacter pylori Eradication for Cronkhite-Canada Syndrome. *Case Rep Gastroenterol*. 2013 Jan;7(1):140-6.
211. Fujii Y, Abe M, Higuchi T, Mizuno M, Suzuki H, Matsumoto S, Ito M, Maruyama N, Okada K, Soma M. The dipeptidyl peptidase-4 inhibitor alogliptin improves glycemic control in type 2

- diabetic patients undergoing hemodialysis. *Expert Opin Pharmacother*. 2013 Feb;14(3):259-67
212. Fujita T, Hemmi S, Kajiwara M, Yabuki M, Fuke Y, Satomura A, Soma M. Complement-mediated chronic inflammation is associated with diabetic microvascular complication. *Diabetes Metab Res Rev*. 2013 Mar;29(3):220-6.
213. Abe M, Okada K, Soma M. Mineral metabolic abnormalities and mortality in dialysis patients. *Nutrients*. 2013 Mar 22;5(3):1002-23.
214. Maruyama N, Abe M, Okada K, Soma M. Changes to prothrombin international normalized ratio in patients receiving hemodialysis. *Int J Clin Pharmacol Ther*. 2013 Apr;51(4):283-7.
215. Kato K, Sugitani M, Matsukawa Y, Sato H, Moriyama M, Soma M. Long-term gastric plasmacytoma follow-up after *Helicobacter pylori* eradication. *Gastrointest Endosc*. 2013 Apr;77(4):674-5.
216. Takagi K, Fujiwara K, Takayama T, Mamiya T, Soma M, Nagase H. DNA hypermethylation of zygote arrest 1 (ZAR1) in hepatitis C virus positive related hepatocellular carcinoma. *Springerplus*. 2013 Apr 10;2(1):150.
217. Shimizu C, Fujita T, Fuke Y, Yabuki M, Kajiwara M, Hemmi S, Satomura A, Soma M. Effects of cyclosporine on bone mineral density in patients with glucocorticoid-dependent nephrotic syndrome in remission. *Int Urol Nephrol*. 2013 Jun;45(3):803-8.
218. Pan S, Nakayama T, Sato N, Izumi Y, Soma M, Aoi N, Ma Y, Hinohara S, Doba N. A Haplotype of the GOSR2 Gene Is Associated with Myocardial Infarction in Japanese Men. *Genet Test Mol Biomarkers*. 2013 Jun;17(6):481-8.
219. Suzuki H, Okada K, Abe M, Maruyama N, Yoshida Y, Baba S, Takashima H, Soma M. Aliskiren reduces home blood pressure and albuminuria in patients with hypertensive

- nephrosclerosis. *Clin Exp Nephrol*. 2013 Jun;17(3):386-95.
220. Pan S, Nakayama T, Sato N, Izumi Y, Soma M, Aoi N, Ma Y. A haplotype of the GOSR2 gene is associated with essential hypertension in Japanese men. *Clin Biochem*. 2013 Jun;46(9):760-5.
221. Abe M, Suzuki H, Okada K, Maruyama N, Inoshita A, Baba S, Takashima H, Soma M. Efficacy analysis of the renoprotective effects of aliskiren in hypertensive patients with chronic kidney disease. *Heart Vessels*. 2013 Jul;28(4):442-52.
222. Abe M, Maruyama N, Suzuki H, Inoshita A, Yoshida Y, Okada K, Soma M. L/N-type calcium channel blocker cilnidipine reduces plasma aldosterone, albuminuria, and urinary liver-type fatty acid binding protein in patients with chronic kidney disease. *Heart Vessels*. 2013 Jul;28(4):480-9.
223. Abe M, Okada K, Suzuki H, Yoshida Y, Soma M. T/L-type calcium channel blocker reduces the composite ranking of relative risk according to new KDIGO guidelines in patients with chronic kidney disease. *BMC Nephrol*. 2013 Jul 1;14(1):135.
224. 阿部雅紀, 丸山範晃, 岡村雅広, 森内正理, 根岸英理子, 古川哲也, 馬場晴志郎, 岡田一義, 相馬正義. 血液透析患者におけるインスリングルリジンの有用性と安全性の検討. *新薬と臨牀*. 2013 Jul:1425-1429.
225. Abe M, Okada K, Soma M. T-type Ca Channel Blockers in Patients with Chronic Kidney Disease in Clinical Practice. *Curr Hypertens Rev*. 2013 Aug;9(3):202-9.
226. Zhou X, Fukuda N, Matsuda H, Endo M, Wang X, Saito K, Ueno T, Matsumoto T, Matsumoto K, Soma M, Kobayashi N, Nishiyama A. Complement 3 activates the renal renin-angiotensin system by induction of epithelial-to-mesenchymal transition of the nephrotubulus in mice. *Am J Physiol Renal Physiol*. 2013 Oct 1;305(7):F957-67.

227. Kamide K, Asayama K, Katsuya T, Ohkubo T, Hirose T, Inoue R, Metoki H, Kikuya M, Obara T, Hanada H, Thijs L, Kuznetsova T, Noguchi Y, Sugimoto K, Ohishi M, Morimoto S, Nakahashi T, Takiuchi S, Ishimitsu T, Tsuchihashi T, Soma M, Higaki J, Matsuura H, Shinagawa T, Sasaguri T, Miki T, Takeda K, Shimamoto K, Ueno M, Hosomi N, Kato J, Komai N, Kojima S, Sase K, Miyata T, Tomoike H, Kawano Y, Ogihara T, Rakugi H, Staessen JA, Imai Y; GEANE study group; HOMED-BP study group. Genome-wide response to antihypertensive medication using home blood pressure measurements: a pilot study nested within the HOMED-BP study. *Pharmacogenomics*. 2013 Nov;14(14):1709-21.
228. Sato A, Nagase H, Obinata D, Fujiwara K, Fukuda N, Soma M, Yamaguchi K, Kawata N, Takahashi S. Inhibition of MMP-9 using a pyrrole-imidazole polyamide reduces cell invasion in renal cell carcinoma. *Int J Oncol*. 2013 Nov;43(5):1441-6.
229. 藤井由季, 阿部雅紀, 樋口輝美, 鈴木紘子, 松本史郎, 水野真理, 菊池史, 岡田一義, 相馬正義. 糖尿病血液透析患者におけるアログリプチンの有効性と安全性の検討. *透析会誌* 46(7):2013 633-640, 2013 Dec
230. Okada K, Abe M, Takashima H, Baba S, Taniguchi M, Suzuki H, Yoshida Y, Oikawa O, Maruyama N, Soma M. Randomized trial of frequent low-efficiency and short hemodialysis/hemofiltration in hemodialysis patients with acute brain injury. *Int J Artif Organs*. 2013 Dec 12;36(11):793-802.
231. Fuke Y, Murata Y, Hemmi S, Kajiwara M, Yabuki M, Fujita T, Soma M. Secondary focal segmental glomerulosclerosis in an adolescent born with a very low birth weight. *Intern Med*. 2014;53(19):2233-6.
232. Fujita T, Hemmi S, Kajiwara M, Yabuki M, Fuke Y, Satomura A, Soma M. Diminished complement-activating capacity through the classical pathway in sera from type 2 diabetes mellitus. *Immunol Invest*. 2014;43(2):170-81.

233. Taniguchi M, Fujiwara K, Nakai Y, Ozaki T, Koshikawa N, Toshio K, Kataba M, Oguni A, Matsuda H, Yoshida Y, Tokuhashi Y, Fukuda N, Ueno T, Soma M, Nagase H. Inhibition of malignant phenotypes of human osteosarcoma cells by a gene silencer, a pyrrole-imidazole polyamide, which targets an E-box motif. *FEBS Open Bio*. 2014 Mar 13;4:328-34.
234. Ikeda K, Fukuda N, Ueno T, Endo M, Kobayashi N, Soma M, Matsumoto K. Role of complement 3a in the growth of mesangial cells from stroke-prone spontaneously hypertensive rats. *Clin Exp Hypertens*. 2014;36(1):58-63.
235. Kojima T, Wang X, Fujiwara K, Osaka S, Yoshida Y, Osaka E, Taniguchi M, Ueno T, Fukuda N, Soma M, Tokuhashi Y, Nagase H. Inhibition of Human Osteosarcoma Cell Migration and Invasion by a Gene Silencer, Pyrrole-Imidazole Polyamide, Targeted at the Human MMP9 NF- κ B Binding Site. *Biol Pharm Bull*. 2014;37(9):1460-5.
236. Tsunemi A, Ueno T, Fukuda N, Watanabe T, Tahira K, Haketa A, Hatanaka Y, Tanaka S, Matsumoto T, Matsumoto Y, Nagase H, Soma M. A novel gene regulator, pyrrole-imidazole polyamide targeting ABCA1 gene increases cholesterol efflux from macrophages and plasma HDL concentration. *J Mol Med (Berl)*. 2014 May;92(5):509-21.
237. Uekusa S, Kawashima H1, Sugito K, Yoshizawa S, Shinojima Y, Igarashi J, Ghosh S, Wang X, Fujiwara K, Ikeda T, Koshinaga T, Soma M, Nagase H. Nr4a3, a possible oncogenic factor for neuroblastoma associated with CpGi methylation within the third exon. *nt J Oncol*. 2014 May;44(5):1669-77.
238. Suzuki R, Fukuda N, Katakawa M, Tsunemi A, Tahira Y, Matsumoto T, Ueno T, Soma M. Effects of an Angiotensin II Receptor Blocker on the Impaired Function of Endothelial Progenitor Cells in Patients With Essential Hypertension. *Am J Hypertens*. 2014 May;27(5):695-701.

239. Kato K, Ohkusa T, Terao S, Chiba T, Murakami K, Yanaka A, Uehara T, Ishii Y, Soma M, Tajiri H. Adjunct antibiotic combination therapy for steroid-refractory or -dependent ulcerative colitis: an open-label multicentre study. *Aliment Pharmacol Ther.* 2014 May;39(9):949-56.
240. Uchida S, Takahashi M, Sugawara M, Saito T, Nakai K, Fujita M, Mochizuki K, Shin I, Morita T, Hikita T, Itakura H, Takahashi Y, Mizuno S, Ohno Y, Ito K, Ito T, Soma M. Effects of the N/L-Type Calcium Channel Blocker Cilnidipine on Nephropathy and Uric Acid Metabolism in Hypertensive Patients With Chronic Kidney Disease (J-CIRCLE Study). *J Clin Hypertens (Greenwich).* 2014 Oct;16(10):746-53.
241. Obinata D, Ito A, Fujiwara K, Takayama KI, Ashikari D, Murata Y, Yamaguchi K, Urano T, Fujimura T, Fukuda N, Soma M, Watanabe T, Nagase H, Inoue S, Takahashi S. Pyrrole-imidazole polyamide targeted to break fusion sites in TMPRSS2 and ERG gene fusion represses prostate tumor growth. *Cancer Sci.* 2014 Oct;105(10):1272-8.
242. Suzuki K, Hayashi Y, Otsuka H, Hashimoto K, Kasamaki Y, Yanai M, Kato K, Nagai Y, Soma M. [Invasive pneumococcal disease of serotype 6C with skin and soft tissue infections in an immunocompetent adult]. *Kansenshogaku Zasshi.* 2014 Nov;88(6):849-54. Japanese.
243. Akita M, Suzuki-Karasaki M, Fujiwara K, Nakagawa C, Soma M, Yoshida Y, Ochiai T, Tokuhashi Y, Suzuki-Karasaki Y. Mitochondrial division inhibitor-1 induces mitochondrial hyperfusion and sensitizes human cancer cells to TRAIL-induced apoptosis. *Int J Oncol.* 2014 Nov;45(5):1901-12.
244. Ito M, Abe M, Maruyama T, Suzuki H, Inoshita A, Okada K, Soma M. Pneumatosis cystoides intestinalis and hepatic portal venous gas on peritoneal dialysis. *Clin Nephrol.* 2014 Nov;82(5):347-50.
245. Ando K, Ohtsu H, Uchida S, Kaname S, Arakawa Y, Fujita T; EVALUATE Study Group (Fujita T, Ando K, Seki G, Tojo A, Shimosawa T, Suzuki H, Arakawa Y, Kubota K, Kiuchi T,

- Yamada A, Kaname S, Kimura K, Uchida S, Yamaguchi T, Ohtsu H, Fukuda N, Kaifu K, Suzuki D, Seki G, Tojo A, Shimosawa T, Takahashi K, Isshiki M, Uchida S, Fukuda N, Soma M, Kaifu K, Suzuki D, Kaname S, Shibagaki Y, Miyashita Y, Umemura S, Kanaoka T, Kouichi K, Ogasawara Y, Inoue S, Kamata K, Tsumura H, Saiki T, Kuwahara Y, Ozono R, Omura M, Katsuya T, Miyairi M, Okuda T, Sato A, Kawana M, Tatumi F, Aihara K, Murakami M, Yasuda G, Arai M, Koba S, Sugiura K, Ikeda H, Ryuuzaki M, Kazama H, Hinoshita F, Onaka H, Ishimitsu T, Iwaida Y, Nomura S, Sawayama Y, Okamoto T, Iyori M, Ota A, Shinoda A, Kawamura M, Kanno K, Kusaka M, Terada Y, Ando S, Ikeda K, Muneyuki T, Inishi Y, Ebihara Y, Hidaka S, Takahashi T, Tsuruoka S, Morita T, Nakano T, Samejima K.). Anti-albuminuric effect of the aldosterone blocker eplerenone in non-diabetic hypertensive patients with albuminuria: a double-blind, randomised, placebo-controlled trial. *Lancet Diabetes Endocrinol.* 2014 Dec;2(12):944-53.
246. Higuchi T, Abe M, Yamazaki T, Mizuno M, Okawa E, Ando H, Oikawa O, Okada K, Kikuchi F, Soma M. Effects of levocarnitine on brachial-ankle pulse wave velocity in hemodialysis patients: a randomized controlled trial. *Nutrients.* 2014 Dec 22;6(12):5992-6004.
247. Igarashi J, Fukuda N, Inoue T, Nakai S, Saito K, Fujiwara K, Matsuda H, Ueno T, Matsumoto Y, Watanabe T, Nagase H, Bando T, Sugiyama H, Itoh T, Soma M. Preclinical Study of Novel Gene Silencer Pyrrole-Imidazole Polyamide Targeting Human TGF- β 1 Promoter for Hypertrophic Scars in a Common Marmoset Primate Model. *PLoS One.* 2015 May 4;10(5):e0125295. doi: 10.1371/journal.pone.0125295. eCollection 2015.
248. Shimizu C, Matsumoto K, Fujita T, Fuke Y, Soma M. Imbalance of interleukin-18 and interleukin-18 binding protein in patients with IgA nephropathy implicating renal vasculopathy. *Clin Lab.* 2015;61(1-2):23-30.
249. Fujiwara K, Ghosh S, Liang P, Morien E, Soma M, Nagase H. Genome-wide screening of aberrant DNA methylation which associated with gene expression in mouse skin cancers. *Mol Carcinog.* 2015 Mar;54(3):178-88.

250. Abe M, Oikawa O, Okada K, Soma M. Urinary angiotensin-converting enzyme 2 increases in diabetic nephropathy by angiotensin II type 1 receptor blocker olmesartan. *J Renin Angiotensin Aldosterone Syst.* 2015 Mar;16(1):159-64.
251. Mishra R, Watanabe T, Kimura MT, Koshikawa N, Ikeda M, Uekusa S, Kawashima H, Wang X, Igarashi J, Choudhury D, Grandori C, Kemp CJ, Ohira M, Verma NK, Kobayashi Y, Takeuchi J, Koshinaga T, Nemoto N, Fukuda N, Soma M, Kusafuka T, Fujiwara K, Nagase H. Identification of a novel E-box binding pyrrole-imidazole polyamide inhibiting MYC-driven cell proliferation. *Cancer Sci.* 2015 Apr;106(4):421-9.
252. Maruyama T, Fukuda N, Matsumoto T, Kano K, Endo M, Kazama M, Kazama T, Ikeda J, Matsuda H, Ueno T, Abe M, Okada K, Soma M, Matsumoto K, Kawachi H. Systematic implantation of dedifferentiated fat cells ameliorated monoclonal antibody 1-22-3-induced glomerulonephritis by immunosuppression with increases in TNF-stimulated gene 6. *Stem Cell Res Ther.* 2015 Apr 16;6(1):80.
253. Kobayashi H, Abe M, Okada K, Tei R, Maruyama N, Kikuchi F, Higuchi T, Soma M. Oral zinc supplementation reduces the erythropoietin responsiveness index in patients on hemodialysis. *Nutrients.* 2015 May 15;7(5):3783-95
254. Hemmi S, Matsumoto N, Jike T, Obana Y, Nakanishi Y, Soma M, Hemmi A. Proximal tubule morphology in rats with renal congestion: a study involving the in vivo cryotechnique. *Med Mol Morphol.* 2015 Jun;48(2):92-103.
255. Yoshizawa S, Fujiwara K, Sugito K, Uekusa S, Kawashima H, Hoshi R, Watanabe Y, Hirano T, Furuya T, Masuko T, Ueno T, Fukuda N, Soma M, Ozaki T, Koshinaga T, Nagase H. Pyrrole-imidazole polyamide-mediated silencing of KCNQ1OT1 expression induces cell death in Wilms' tumor cells. *Int J Oncol.* 2015 Jul;47(1):115-21.

256. Suzuki-Karasaki Y, Fujiwara K, Saito K, Suzuki-Karasaki M, Ochiai T, Soma M. Distinct effects of TRAIL on the mitochondrial network in human cancer cells and normal cells: role of plasma membrane depolarization. *Oncotarget*. 2015 Aug 28;6(25):21572-88.
257. Abe M, Maruyama N, Oikawa O, Maruyama T, Okada K, Soma M. Urinary ACE2 is associated with urinary L-FABP and albuminuria in patients with chronic kidney disease. *Scand J Clin Lab Invest*. 2015 Sep;75(5):421-7.
258. Saito K, Fukuda N, Shinohara K, Masuhiro Y, Hanazawa S, Matsuda H, Fujiwara K, Ueno T, Soma M. Modulation of the EMT/MET process by pyrrole-imidazole polyamide targeting human transforming growth factor- β 1. *Int J Biochem Cell Biol*. 2015 Sep;66:112-20.
259. Kobayashi H, Haketa A, Abe M, Tahira K, Hatanaka Y, Tanaka S, Ueno T, Soma M. Unusual Manifestation of Graves' Disease: Ventricular Fibrillation. *Eur Thyroid J*. 2015 Sep;4(3):207-12.
260. Sezai A, Soma M, Nakata KI, Osaka S, Ishii Y, Yaoita H, Hata H, Shiono M. Comparison of febuxostat and allopurinol for hyperuricemia in cardiac surgery patients with chronic kidney disease (NU-FLASH trial for CKD). *J Cardiol*. 2015 Oct;66(4):298-303.
261. Ozaki Y, Fujiwara K, Ikeda M, Ozaki T, Terui T, Soma M, Inazawa J, Nagase H. The oncogenic role of GASC1 in chemically induced mouse skin cancer. *Mamm Genome*. 2015 Dec;26(11-12):591-7.
262. Abe M, Maruyama N, Maruyama T, Okada K, Soma M. A Trial of Pitavastatin Versus Rosuvastatin for Dyslipidemia in Chronic Kidney Disease. *J Atheroscler Thromb*. 2015 Dec 1;22(12):1235-47
263. Abe M, Soma M. Multifunctional L/N- and L/T-type calcium channel blockers for kidney

protection. *Hypertens Res.* 2015 Dec;38(12):804-6.

264. Inami M, Fukushima A, Ueno T, Yamada T, Tsunemi A, Matsumoto Y, Fukuda N, Soma M, Moriyama M. Reduction of Dimethylnitrosamine-Induced Liver Fibrosis by the Novel Gene Regulator PI Polyamide Targeting Transforming Growth Factor β 1 Gene. *Biol Pharm Bull.* 2015 Dec 1;38(12):1836-42.
265. Fujita T, Watanabe H, Murata Y, Hemmi S, Yabuki M, Fuke Y, Satomura A, Soma M. Plasma C1q/tumor necrosis factor-related protein 9; a promising biomarker for diabetic renal vascular injury. *Minerva Urol Nefrol.* 2015 Dec 3.
266. Kobayashi H, Haketa A, Ueno T, Suzuki R, Aoi N, Ikeda Y, Tahira K, Hatanaka Y, Tanaka S, Otsuka H, Abe M, Fukuda N, Soma M. Subtype Prediction in Primary Aldosteronism: Measurement of Circadian Variation of Adrenocortical Hormones and 24-h Urinary Aldosterone. *Clin Endocrinol (Oxf).* 2015 Dec 13.
267. Abe M, Okada K, Maruyama N, Takashima H, Oikawa O, Soma M. Comparison of Clinical Trajectories before Initiation of Renal Replacement Therapy between Diabetic Nephropathy and Nephrosclerosis on the KDIGO Guidelines Heat Map. *J Diabetes Res.* 2016;2016:5374746.
268. Higuchi T, Abe M, Yamazaki T, Okawa E, Ando H, Hotta S, Oikawa O, Kikuchi F, Okada K, Soma M. Levocarnitine Improves Cardiac Function in Hemodialysis Patients With Left Ventricular Hypertrophy: A Randomized Controlled Trial. *Am J Kidney Dis.* 2016 Feb;67(2):260-70.
269. Saito K, Asai T, Fujiwara K, Sahara J, Koguchi H, Fukuda N, Suzuki-Karasaki M, Soma M, Suzuki-Karasaki Y. Tumor-selective mitochondrial network collapse induced by atmospheric gas plasma-activated medium. *Oncotarget.* 2016 Mar 3.

270. Yanai M, Gon Y, Suzuki K, Hayashi Y, Hasumi S, Otsuka H, Soma M. Clinical usefulness of serum 2'-5'-oligoadenylate synthetase for early diagnosis of viral infections among febrile adult patients. *Infect Dis (Lond)*. 2016 May;48(5):338-42.

症例報告

1. 相馬正義、本多正信、波多野道信：Cushing Syndrome を呈した副腎癌の1例について。第1回 op'-DDD 研究会議事録：pp.60-66, 1980.2
2. 植田康久、本多正信、藤岡孝之、相馬正義、永嶋康夫、井上毅、土屋真、泉洋一、白土寿、波多野道信：特発性アルドステロン症の1例。日大医学雑誌 41:pp321-327, 1982.3
3. 相馬正義、本多正信、波多野道信：Cushing 症候群を呈した副腎皮質癌の1症例。ホルモンと臨床 '82 増刊号:pp201-205, 1982.
4. 渡辺吉康、本多正信、水野学、井上毅、岸均、泉洋一、真鍋亮子、渡部秀人、永嶋康夫、相馬正義、白土寿、波多野道信、佐々英一：著名な塩類喪失を示した下垂体機能低下症の一例。ホルモンと臨床 '83 増刊号:pp30-35, 1983
5. 湊昌也、本多正信、渡辺昌司、福田昇、相馬正義、永嶋康夫、泉洋一、白土寿、波多野道信：内分泌学的異常、Adams-Stokes 様発作、心膜液貯留、脳萎縮を伴った神経性食欲不振症の一男性例。日本内科学会雑誌 77:pp.1417-1420, 1988.9
6. 相馬正義、本多正信、湊昌也、渡辺昌司、福田昇、永嶋康夫、泉洋一、白土寿、波多野道信：バーター症候群様病態を呈した3症例。ホルモンと臨床 11:pp.217-220, 1989. '89 春季増刊号
7. 笠巻祐二、松本太郎、小松一俊、相馬正義、栃原敏彦、奈倉勇爾、小沢友紀雄、波多野道信、沼口満津男：高齢で腹部大動脈瘤破裂をきたした1例。日本臨床内科医学会誌 4:63-65, 1990.
8. 奈尾俊子、相馬正義、渡辺昌司、渡辺吉康、泉洋一、波多野道信：慢性甲状腺炎を合併したACTH単独欠損症の一例。ACTH RELATED PEPTIDS 2:pp.96-101, 1991.2
9. 伊藤みき、泉洋一、相馬正義、八杉忠夫：プロラクチン産生腫瘍にGraves

病、I g A腎症をともなった症例。ホルモンと臨床 40 卷秋季増刊号 内分泌興味ある症例 第19集 : pp.93-98, 1992.4

10. 阿部好伸、相馬正義、松岡昌子、中山智祥、伊藤みき、渡辺昌司、渡辺吉康、泉洋一、八杉忠夫：Cushing 症候群を呈した副腎腺腫として手術され約10年後に再発した副腎癌の一例。ホルモンと臨床 40 卷秋季増刊号 内分泌興味ある症例 20集 : pp.141-144, 1992.10
11. 中山智祥、相馬正義、久保淳、松岡昌子、阿部好伸、伊藤みき、渡辺昌司、渡辺吉康、泉洋一、八杉忠夫：妊娠合併のCushing 症候群-自験例副腎腺腫1例と文献的考察-。日本内分泌学会誌 68:pp.1130-1149, 1992.10
12. 宋滋章、磯村政保、渡辺太次郎、飯島隆紀、田中直英、相沢敏晴、伊藤和郎、岩崎有良、松尾裕、相馬正義、泉洋一、岡野国雄：ACTH単独欠損症の一例。日大医学雑誌 52:pp35-42, 1993.1
13. 細川留美、相馬正義、鳴瀬裕爾、渡辺昌司、森内正人、大井洋之、上松瀬勝男：Empty sella を伴うACTH単独欠損症の一例。日大医学雑誌 55:95-100, 1996.2
14. 中山麻里、福田昇、渡辺吉康、相馬正義、泉洋一、上松瀬勝男：オクトレオチドが奏功したネルソン症候群の1例。ホルモンと臨床 '96 夏期増刊号 内分泌興味ある症例第28集 pp101-103, 1996.8
15. 高橋雪絵、福田昇、相馬正義、泉洋一、上松瀬勝男：内分泌異常を伴いオクトレオチドが奏功した線維性骨異形成の1例。ホルモンと臨床 '96 夏期増刊号 内分泌興味ある症例 第28集 pp101-103, 1996.8
16. Nakayama T, Asai Y, Soma M, Aoki K, Tobe H, Ozawa Y, Kanmatsuse K, Nezu T, Amano S. A family with multiple endocrine neoplasia type 2A. Nihon Univ J Med 41'309-318, 1999.10
17. Nakayama T, Nakayama K, Takahashi Y, Ohkubo K, Tobe H, Soma M, Ozawa Y, Kanmatsuse K, Nakamura M, Hironaga T, Makizumi Y, Nagura H: Case of spinocerebellar

ataxia type 1 showing high intensity lesions in the frontal white matter on T2-weighted magnetic resonance images. Med Sci Monit 7(2):299-303, 2001. 4

18. 天野定雄、久保井洋一、斉藤良人、西尾知、大原守貴、根津健、三宅洋、大井田尚継、秦怜志、福沢正洋、相馬正義、上松瀬勝男：内視鏡補助下甲状腺部分切除および副甲状腺摘出術の経験。日大医学雑誌 60:144-148, 2001.3
19. 中山智祥、山口賢二、及川 治、相馬正義、泉 洋一、小沢友紀雄、上松瀬勝男、柚木隆寛、平野大作、高橋雪絵、斎藤友昭：副腎腺腫によるクッシング症候群の一例。板橋区医師会医学会誌 5:pp.201~203, 2001.5
20. 松崎真和、相馬正義、佐藤三佳乃、鈴木亮、赤羽正史、岡田一義、松本紘一、上松瀬勝男、岡田清巳、石田肇、平野大作、杉谷雅彦：ACTH非依存性両側副腎皮質大結節性過形成（AIMAH）の一例。日大医学雑誌 62(2):121-125, 2003
21. 山崎俊男、樋口輝美、大西貞彦、松本史郎、矢吹美奈子、北島尚子、及川治、丸山範晃、羽木千尋、岡田一義、福田昇、藤田宣是、相馬正義、松本紘一：Hypereosinophiliaを呈した透析困難症の一例。日大医学雑誌 64(3):160-163,2005
22. Sato M, Soma M, Nakayama T, Kosuge K, Suzuki R, Okada K, Komatsu K, Sugitani M, Matsumoto K. A case of adrenocorticotropin-independent bilateral adrenal macronodular hyperplasia (AIMAH) with primary hyperparathyroidism. Endocrine Journal 2006, 53(1):111-117.
23. Higuchi T, Yamazaki T, Ohnishi Y, Matsumoto S, Yabuki M, Kitajima S, Toshida M, Maruyama Y, Oikawa O, Okada K, Fukuda N, Soma M, Matsumoto K. A case report of hemodialysis intolerance with eosinophilia. Ther Apher Dial. 2007 Feb;11(1):70-3.
24. 降籟隆二、平野亘、河合真紀子、久保英之、内山真、小菅琴子、田平佳子、伊田則子、相馬正義、松本紘一：悪性症候群を呈したACTH単独欠損症を伴った統合失調症の一例。日大医学雑誌 66(1):110-115,2007.2
25. Obinata D, Yamaguchi K, Hirano D, Yoshida T, Soma M, and Takahashi S. Preoperative

management of Cushing's syndrome with metyrapone for severe psychiatric disturbances. *International Journal of Urology* 2008 Apr;15(4):361-2.

26. 岡田一義, 阿部雅紀, 水盛邦彦, 小林伸一郎, 佐々木裕和, 鈴木紘子, 谷口真知子, 逸見聖一朗, 北井真貴, 鈴木 緑, 奈倉千苗美, 池田和也, 梶原真実子, 清水千枝, 矢吹美奈子, 松本史郎, 伊藤 謙, 井下篤司, 丸山高史, 福家吉伸, 藤田宜是, 相馬正義. 頭蓋内圧亢進症急性期に高張グリセロール併用頻回低効率短時間血液透析 (frequent, low-efficient and short hemodialysis : FLESHD) が有用であった維持血液透析症例. *透析会誌*. 2011;44(6):581-587
27. 丸山高史, 阿部雅紀, 奈倉千苗美, 鈴木 緑, 岡田一義, 相馬正義. 心筋症による重症心不全に腹膜透析を導入した 1 例. *臨床透析*. 2011;27(5):585-591.
28. Suzaki A, Hayashi K, Kosuge K, Soma M, Hayakawa S. Disseminated gonococcal infection in Japan: a case report and literature review. *Intern Med*. 2011;50(18):2039-43.
29. 高野智圭, 熊澤文雄, 野田彰浩, 野村舟三, 高橋恵子, 菅谷健一, 小山大輔, 新谷榮崇, 鈴木紘子, 古川 誠, 岡本直樹, 大木隆史, 阿部雅紀, 林 伸一, 小林朋子, 岡田和義, 高橋典明, 木下浩作, 相馬正義, 丹正勝久, 橋本 修. ステロイドパルス療法, エンドトキシン吸着療法およびシクロスポリン A が著効した急性間質性肺炎の一例. *日大医誌*. 2012. 71(6):455-459
30. Abe M, Okada K, Maruyama N, Matsumoto S, Fuke Y, Fujita T, Soma M, Matsumoto K. A case of femoral hemorrhage in a patient with microscopic polyangiitis with low levels of myeloperoxidase-antineutrophil cytoplasmic autoantibody. *Clin Exp Nephrol*. 2011 Jun;15(3):414-8.
31. Fuke Y, Hemmi S, Kajiwara M, Yabuki M, Fujita T, Soma M. Oligomeganephronia in an adult without end stage renal failure. *Clin Exp Nephrol*. 2012 Apr;16(2):325-8.
32. Suzuki H, Fujita T, Fuke Y, Yabuki M, Kajiwara M, Ishihara Y, Hemmi S, Soma M. A case of ANCA-associated vasculitis with glomerular eosinophilic infiltration: a possible pathogenic implication. *Mod Rheumatol*. 2013 May;23(3):564-71

33. Suzuki H, Abe M, Tahira K, Ito M, Takashima H, Baba S, Okada K, Soma M. Successful treatment of pheochromocytoma in a patient with hemodialysis: a case report and review of the literature. *Ren Fail*. 2013 Nov;35(10):1429-33.

34. 鈴木清澄, 林 悠太, 大塚博雅, 橋本賢一, 笠巻祐二, 矢内 充, 加藤公敏, 永井 悠, 相馬正義. 健常成人に皮膚軟部組織感染症で発症した莢膜型 6C による侵襲性肺炎球菌感染症の 1 例. *感染症学雑誌*, 2014 Nov;88(6):849-853.

総説

1. 相馬正義、本多正信、泉洋一、波多野道信： Hypertensinogenic steroid. 日本臨床 47:pp.2104-2108, 1989.9
2. Soma M, Nakayama T, Kanmatsuse K. NOS gene polymorphism and its influence on cardiovascular disease. Cur Opinion in Nephrol Hypertens 8:83-87, 1999.1
3. 相馬正義：どのような場合に二次性高血圧症の精査をすべきか。Medical Practice 17:783-786, 2000.5
4. 相馬正義：遺伝子と健康。教育医療 28(7):4-5, 2002.7
5. 相馬正義：加齢とともに起こる骨粗鬆症—治療はここまで可能—。教育医療 29(10):2-3, 2003.10
6. 相馬正義：糖尿病と β 遮断薬の関係は？肥満と糖尿病 3(1):144-145, 2004.1
7. 相馬正義：糖尿病合併高血圧患者にCa拮抗薬の第1選択は適切か？Medicina 41(1):123-124, 2004.1
8. 相馬正義：高血圧と遺伝子異常。日大医学雑誌 63(11,12):527-528, 2004.
9. 相馬正義：候補遺伝子アプローチ（日大方式）。血圧 12(8):839-842, 2005.
10. 相馬正義：PBLテュトリアルシステム。日大医学雑誌 64(6)337-339, 2005
11. 阿部雅紀、岡田一義、松本紘一、相馬正義。第8回日本高齢者腎不全研究会「高齢透析患者の糖尿病管理」。腎と透析 69(4)511-515, 2010.10
12. 阿部雅紀、相馬正義。CKD患者における高血圧管理。Angiology Frontier 9(4)347:34-44, 2010.12
13. 阿部雅紀、岡田一義、松本紘一、相馬正義。高齢透析患者の糖尿病管理。腎と透

析. 2010 Oct;69(4):511-515

14. 中山智祥、青井則子、佐藤直之、佐藤三佳乃、小菅琴子、泉洋一、相馬正義、松本絃一。ギテルマン症候群の遺伝学的検査体制の構築。臨床病理（日本臨床検査医学会誌）。2010 Feb 20; 58(2):156-161.
15. 松本絃一、遠藤守人、福家吉伸、伊藤 謙、清水千枝、梶原真実子、丸山高史、井下篤司、阿部雅紀、岡田一義、藤田宜是、福田 昇、相馬正義. ANCA 関連腎炎. 医学と薬学. 2011 Mar;65(3):319-325.
16. Okada K, Abe M, Soma M. Implementation of a cooperative program for peritoneal dialysis. Contrib Nephrol. 2012;177:84-92.
17. 相馬正義. 遺伝子解析の現状と近未来. 東京都医師会雑誌. 2012 Feb; 65(2):4-6
18. 笠巻祐二, 橋本賢一, 相馬正義. 症状時の心電図をどのようにして記録するか? Medicina vol.50 No.13 2013-12:2131-2135
19. 相馬正義. 高血圧合併時の薬剤投与: 注意と禁忌薬. 成人病と生活習慣病. 2014 vol.44, No2:152-157.

著書

1. Matsuoka M, Watanabe Y, Soma M, Izumi Y, Yasugi T: Effects of docosahexaenoic acid on blood pressure, fatty acid composition of kidney and renal prostaglandins in Dahl salt sensitive rats. *Advances in Polyunsaturated Fatty Acid Research*, edited by T. Yasugi, H. Nakamura and M. Soma, pp183-184, Excerpta Medica, Amsterdam, 1993.5
2. Watanabe Y, Matsuoka M. Soma M, Izumi Y, Yasugi T: Effect of docosahexaenoic acid ethylester on fatty acid composition of aorta and liver in Dahl salt sensitive or resistant rats. *Advances in Polyunsaturated Fatty Acid Research*, edited by T. Yasugi, H. Nakamura and M. Soma, pp185-186, Excerpta Medica, Amsterdam, 1993.5
3. 相馬正義、泉洋一、上松瀬勝男：高血圧ケースセミナー。矢崎義雄監修、メディカルジャーナル社、1997.8
4. Nakayama T, Soma M, Kanmatsuse K. Organization of the human prostacyclin synthase gene and association analysis of a novel CA repeats in essential hypertension. *Recent Advances in Prostaglandin, Thromboxane, and Leukotriene Research*, Edited by Sinzinger et al. pp127-130, 1998.4
5. 相馬正義：研修医必携 薬物療法と禁忌。視床下部・下垂体疾患。岡田一義、福田昇、松本絃一監修、東京医学社、2005.4
6. 相馬正義：研修医必携 薬物療法と禁忌。骨粗鬆症。岡田一義、福田昇、松本絃一監修、東京医学社、2005.4
7. 相馬正義：新・病態生理でできた内科学 4. 内分泌疾患。村川裕二総監修、医学教育出版、2006.9
8. 相馬正義：より良い診療をめざして他診療科との連携—甲状腺機能障害。金原出版眼科 52(6):799-05, 2010.6.5
9. 相馬正義：高血圧。東京医学社 成人病と生活習慣病（日本成人病（生活習慣病）学会準機関誌） 40(9): 1039-1041, 2010.9.15

10. 相馬正義: CKD における Ca 拮抗薬の抗蛋白尿効果. 先端医学者 血圧 17(11): 78-80, 2010.11.1
11. 阿部雅紀、相馬正義: CKD 患者における高血圧管理. メディカルレビュー社 Angiology Frontier 9(4); 34-44, 2010.12.1
12. 富野康日己、相馬正義 (他 37 名): 医療禁忌ポケットマニュアル (内分泌内科). 医歯薬出版株式会社. 29-35
13. 相馬正義、村川裕二: 新・病態生理でできた内科学〈4〉内分泌疾患. 医学教育出版社. 2012.10
- 14.

その他

1. 相馬正義。学会トピックス。第25回 日本高血圧学会総会。Atherothrombosis 6(1):39, 2004
2. 岡田一義、工藤たみよ、小松一俊、相馬正義、斉藤聡、松本紘一。インフルエンザ感染防止対策マニュアルの問題点。日大医学雑誌、63(1):43-45, 2004.
3. 相馬正義、松本紘一、道場信孝、日野原重明。ビタミンD受容体遺伝子多型は高齢者の腰椎骨密度と相関する。財団法人ライフ・プランニングセンター研究業績年報、2004(25):58-59, 2005
4. 相馬正義。生活習慣病の解明、DNA解析が予防と治療に寄与。日大板橋病院だより、No.21,pp11-13, 2005.7
5. 相馬正義。日本大学医学部がめざすPBLテュートリアル。日大医学雑誌、65(1):79-80,2006
6. 相馬正義。カロリー制限と寿命と遺伝子。東京都医師会雑誌、第60巻 第6号(19.7.15)
7. 心筋梗塞の遺伝的要因を判定するための方法及びこれにしようされるオリゴヌクレオチド (特願 2000-336676)
8. 循環器疾患の遺伝的要因を判定するための方法及びこれに使用されるオリゴヌクレオチド (特願 2000-167331)
9. 本態性高血圧症の判定方法 (特願 2005-244730)
10. 脳梗塞の判定方法 (特願 2006-010431)
11. 相馬正義。医師臨床研修制度の見直し。東京都医師会雑誌、2009 Feb;62(2):4-5.

12. 相馬正義、松本紘一、道場信孝、日野原重明：ビタミンD受容体遺伝子多型は高齢者の腰椎骨密度と相関する．財団法人ライフ・プランニング・センター研究業績年報—2004, 第25号. 2005 Oct; 25: 58-59
13. 相馬正義：44-46
14. 権 寧博, 矢内 充, 相馬正義. 抗インフルエンザ薬. 日大医学雑誌. 2011;70(5):273
15. 岡田一義, 阿部雅紀, 相馬正義. PDによるQOL向上を目指して. 臨床透析. 2011;27(10):
16. 藤田之彦, 橋本 修, 竹内 仁, 相馬正義, 亀井 聡, 相澤 信, 片山容一. 日本大学医学部における26年間の医学教育セミナーとワークショップを振り返って. 日大医学雑誌. 2012.Jun;71(3): 184-189
17. 鈴木清澄, 権 寧博, 大塚博雅, 林 悠太, 蓮見禎行, 須崎 愛, 山中弥太郎, 矢内 充, 高橋 悟, 相馬正義. 海綿腎. 日大医誌. 2012.Dec;71(6):365-366
18. 林 悠太, 鈴木清澄, 蓮見禎行, 大塚博雅, 須崎 愛, 権 寧博, 矢内 充, 相馬正義. Crowned Dens Syndrome. 日大医学雑誌. 2013.Feb;72(1):2-3.
19. 相馬正義. 職場における腎疾患対策. 東京都医師会・日本大学医師会産業医研修会 講演集. 2013.